

**REPEATED CONFUSIONAL STATES FOLLOWING
DISCONTINUATION OF PROXETINE IN A 51-YEAR-OLD
WOMEN SUFFERING FROM PSYCHOTIC DEPRESSION**

Horst J. Koch

Chefarzt der Klinik für Psychiatrie
Heinrich-Braun-Klinikum Zwickau
Karl-Keil Str. 35, 08060 Zwickau, Germany

Contact: Horst J. Koch MD PhD DCPSA MFPM
Chefarzt der Klinik für Psychiatrie,
Heinrich-Braun-Klinikum Zwickau
Karl-Keil Str. 35, 08060 Zwickau, Germany
Email: horst.koch@hbk-zwickau.de

A 51-year-old woman suffering from depression with psychotic symptoms including a history of meningitis and epilepsy since childhood was treated with paroxetine, olanzapine and lamotrigine for years. In the periods she interrupted paroxetine administration, she developed each time a confusional state requiring intensive psychiatric care. She recovered in a few days after readministration of paroxetine. Clinicians should be aware of severe withdrawal reactions after discontinuation of SSRI, particularly in patients with neurological history. *Acta Medica Medianae* 2015;54(3):51-53.

Key words: *discontinuation syndrome, paroxetine, confusional state, neurological disease*