

Scientific Journal of the Faculty of Medicine in Niš 2010;27(1):13-18

Original article ■

Effects Of Simvastatin Therapy On Bone Mineral Density In Hypercholesterolemic Postmenopausal Woman

Todorka Savić¹, Dimitrije Janković², Irena Janković³, Vladmila Bojanić⁴, Boris Đinđić⁴, Nataša Miladinović-Tasić⁵, Ivan Tasić¹

¹Institute for Treatment and Rehabilitation „Niška Banja“, Serbia, ²Swiss Pharm Serbia, ³Clinic of Plastic Surgery, Clinical Center Niš, Serbia, ⁴Institute of Pathophysiology, Faculty of Medicine in Niš, Serbia, ⁵Public Health Institute Niš, Serbia

SUMMARY

The aim of the study was to evaluate the effects of simvastatin therapy on bone mineral density in hypercholesterolemic postmenopausal women.

Fifty-three women, mean age $62,92 \pm 6,08$ years treated at the Institute for Treatment and Rehabilitation Niška Banja were included into this study. All patients were divided into two groups: I group - 32 women with total cholesterol level ($Hol \geq 7,8$ mmol/l) and II control group - 21 women with total cholesterol less than 7,8 mmol/l. Triglycerides (Tg), total cholesterol (Hol), HDL and LDL cholesterol (HDL-C and LDL-C) were determined. Bone mineral density (BMD) was determined on the lumbar spine (L_1-L_4), by "LUNAR DPX" densitometer. The results were expressed as absolute values (g/cm^2) and T score. All patients were treated with lifestyle and dietary modification, while patients in group I were treated with 20 mg of simvastat per day. Biochemical markers and bone mineral density were determined at the start and 12 months after.

Simvastatin therapy 20mg/day significantly reduced Hol (-12,68%; $p < 0,01$) and LDL-C (-17,99%; $p < 0,005$) levels after 12 months of follow-up. The concentration of TG and HDL-C showed insignificant changes. In II group, there were no significant changes in lipid parameters after 12 months of hypolipemic diet regimen. Increase in BMD +2,81%, (NS) was registered in I group, while decrease in BMD -3,45% (NS) was reported in II group. There was an inverse correlation between BMD changes and LDL-C ($r = -0,08$; $p = 0,686$) in the patients on statin therapy compared to I group ($r = 0,166$, $p = 0,626$). Also, in the same group, an inverse correlation between BMD and Hol ($r = -0,144$; $p = 0,476$) was reported, which was similar to I group ($r = -0,125$; $p = 0,715$). These results did not reach statistical significance.

Conclusion: Simvastatin therapy showed a positive trend and increase in bone mineral density in hypercholesterolemic postmenopausal women after 12 months of follow-up, however, without statistically significant difference.

Key words: bone mineral density, statin therapy, postmenopausal women

Corresponding author:

Todorka Savić •

tel. 064/246 7501 •

e-mail: todorka@medfak.ni.ac.rs •

INTRODUCTION

Statins are hypolipemic drugs with blocking effects on 3-hydroxy 3-methyl glutaryl coenzyme A (HMG-CoA) reductase, the key enzyme in cholesterol synthesis. They have been used in the therapy of hypercholesterolemia as well as for primary and secondary coronary artery disease prevention, with proven efficacy throughout numerous studies (1).

Besides limiting effects on cholesterol synthesis and reduced mevalonate production, they exert some other (pleiotropic) effects apart from their hypolipemic effects. This pleiotropic effects are vasodilatation, antithrombotic, antioxidative, anti-inflammatory and antiproliferative effects (2). The effects of statins on bone mineral metabolism are of huge interest for postmenopausal women with increased risk for coronary artery disease development. This effect is mediated throughout several mechanisms: signaling proteins of osteoclastic activity reduction (3) and increased expression of morphogenetic bone protein 2 gene (4). These mechanisms could be the rationale for statins therapy in osteoporotic women. Some studies have shown decreased risk for bone fractures in hypercholesterolemic postmenopausal women on statin therapy (5-12). There are many questions related to the effects of statin therapy on bone mineral density. Some well-designed and large prospective studies are needed to provide answers on these questions.

AIMS

The aim of the study was to evaluate the effects of simvastatin therapy on bone mineral density in hypercholesterolemic postmenopausal women.

PATIENTS AND METHODOLOGY

Fifty-three women, mean age $62,92 \pm 6,08$ years treated at the Institute for Treatment and Rehabilitation "Niška Banja" in Niška Banja were included into this study. All the patients were hypercholesterolemic postmenopausal women. They were on cardiovascular rehabilitation program prior to study inclusion. The patients with secondary osteoporosis, endocrine or systemic diseases, steroid or hormonal substitution therapy, therapy with thiazides, bisphosphonates or statins were excluded from the study.

All patients were divided into two groups: group I - 32 women with total cholesterol level ($Hol \geq 7,8$ mmol/l) and II control group - 21 women with total cholesterol less than 7,8 mmol/l. All patients had diet with less than 30% of fats in total energy input, less than 7% of saturated fatty acids and less than 200mg/day of cholesterol. They also had therapy with 1200 mg/day of calcium and 400 IJ/day of vitamin D. Patients in group I were treated with 20 mg of simvastat per day.

The blood samples were taken in fasting state. Triglycerides (Tg), total cholesterol (Hol) and HDL cholesterol (HDL-C) were determined by Randox reagents with „Synchron CX5” analyser. LDL cholesterol (LDL-C) was determined by indirect Friedwald method. Bone mineral density (BMD) was determined on the lumbar spine (L_1-L_4), by "LUNAR DPX" densitometer. The results were expressed as absolute values (g/cm^2) and T score (SD deviation from young healthy population referent values). Biochemical markers and bone mineral density were determined at the start and 12 months after.

Statistical analysis was done by SPSS software, using average standard deviation, Student's t-test and Pearson's test for linear correlation.

RESULTS

At the beginning of the study, 81,13% of patients from both groups had decreased bone mineral density (90,62% patients in I group and 66,67% patients in II group). Average BMD was similar in both groups (Table 1).

Mean age and body mass index (BMI) of examined patients did not significantly differ between two groups (Table 1). Plasmatic level of total cholesterol in all patients was $7,66 \pm 1,71$ mmol/l at the beginning of the study ($8,08 \pm 1,74$ mmol/l in patients from I group and $6,6 \pm 1,08$ mmol/l in patients from II group, $p < 0,005$). Average LDL-C concentration was significantly higher in patients from I group compared to II group ($6,12 \pm 1,74$ mmol/l vs. $4,59 \pm 0,99$ mmol/l, $p < 0,001$). Other examined lipid parameters did not show statistical difference between the groups (Table 1).

Simvastatin therapy 20 mg/day significantly reduced Hol levels (-12,68%; $p < 0,01$) and LDL-C (-17,99%; $p < 0,005$) after 12 months of follow-up. Concentration of TG and HDL-C showed insignificant changes of Tg (+6,36%) and HDL-C (+2,22%) compared to starting values. In II group we did not register significant changes in lipid parameters after 12 months of hypolipemic diet regimen. Increasing of BMD +2,81%, (NS) was registered in I group, while decreasing of BMD -3,45% (NS) was observed in II group (Table 2).

Comparison of BMD changes in groups I and II after 12 months of follow-up (Table 3) showed statistically significant difference ($p < 0,0006$).

Patients on statin therapy had inverse correlation between BMD changes and LDL-C ($r = -0,08$; $p = 0,686$) compared to I group ($r = 0,166$, $p = 0,626$). They also had inverse correlation between BMD and Hol ($r = -0,144$; $p = 0,476$) similarly to I group ($r = -0,125$; $p = 0,715$). These results did not reach statistical significance.

Table 1. Investigated parameters at the beginning of the study

Investigated parameters	Group I (n=32)	Group II (n=21)	p
Age (years)	63,11±6,27	62,78±5,01	NS
BMI (kg/m ²)	25,25±2,87	26,73±2,04	NS
BMD (g/cm ²)	0,96±0,11	1,04±0,18	NS
Hol (mmol/l)	8,08±1,74	6,6±1,08	<0,005
TG (mmol/l)	1,75±0,66	1,72±0,58	NS
HDL-C (mmol/l)	1,17±0,24	1,22±0,24	NS
LDL-C (mmol/l)	6,12±1,74	4,59±0,99	<0,001

BMI-body mass index; BMD - bone mineral density; Hol - total cholesterol; Tg- triglycerides; HDL-C - HDL cholesterol; LDL-C - LDL cholesterol

Table 2. Changing of BMD and lipid parameters during the study

Parameters	Group I			Group II		
	beginning	after 12 months	Δ(%)	beginning	after 12 months	Δ(%)
	— X±SD			— X±SD		
BMD (g/cm ²)	0,96±0,11	0,99±0,11	2,81	1,04±0,18	1,01±0,18	-3,45
Hol (mmol/l)	8,08±1,74	7,06±1,14**	-12,68	6,6±1,08	6,44±0,61	-2,41
TG (mmol/l)	1,75±0,66	1,86±0,91	6,36	1,72±0,58	2,01±0,61	16,54
HDL-C (mmol/l)	1,17±0,24	1,19±0,27	2,22	1,22±0,24	1,14±0,26	-6,45
LDL-C (mmol/l)	6,12±1,74	5,02±1,11***	-17,99	4,59±0,99	4,58±0,69	-0,19

p<0,01, *p<0,005

Table 3. Changes of BMD and lipid parameters' values after 12 months in the examined groups

Parameters	Group I		Group II		p value
	X	SD	X	SD	
BMD (g/cm ²)	-0,03	0,04	0,04	0,04	0,0006
Hol (mmol/l)	1,02	1,27	-0,04	0,99	0,011
TG (mmol/l)	-0,11	0,83	-0,28	0,41	0,389
HDL-C (mmol/l)	-0,03	0,31	0,08	0,28	0,313
LDL-C (mmol/l)	1,1	1,27	0,09	0,84	0,004

DISCUSSION

In the past years, some prospective studies showed positive effects of statin therapy on bone mineral metabolism, but none of them aimed to investigate the effects of statins on BMD. Most of them have investigated the association between statin therapy and risk for bone fractures (5-12). Adami et al. (13) consider that lower incidence of bone fractures in patients on statin therapy could be explained by higher BMD observed in patients with higher LDL-C levels. In our study, all women were hypercholesterolemic, while those on statin therapy had higher average Hol and LDL-C levels compared to control group. BMD did not differ among the groups.

In a large retrospective study, Chung et al. demonstrated significant rise of BMD in men with diabetes mellitus type 2 on statin therapy (14). Edwards et al. (15) showed 10% increase of BMD in women on statin therapy compared to dietary regimen without statins. In this study, as well as in the study conducted by Chung et al, different statins were used without analysing particular statin effects on BMD. The effects of fluvastatin and pravastatin on BMD during one-year follow-up were analysed in the study conducted by Watanabe et al. (16). In this study, fluvastatin showed minimal effects on BMD compared to pravastatin which had no effects on BMD. This study showed different effects of statins on cholesterol reduction as well as BMD changes.

Montagnani et al. (17) showed positive effects of simvastatin therapy 40 mg/day on BMD in postmenopausal hypercholesterolemic women compared to normolipidemic women in one-year follow-up study. In this study, the authors did not evaluate the association between lipid profile and BMD considering the results from Rejinmark et al. (18) who showed that statins therapy modulate bone cells function by antiresorptive effects.

Our study indicates the increase in BMD in hypercholesterolemic postmenopausal women on statin therapy compared to women without statins. In our study, the increase in BMD in patients on statin therapy was smaller compared to the results from Montagnani et al (17). This was an expected result as we used simvastatin 20 mg/day. However, the intergroup analysis in our study showed statistically significant change of BMD compared to Montagnani et al¹⁷. BMD in hypercholesterolemic women from control group showed higher decrease compared to results from Montagnani et al. in normolipidemic women (17). This indirectly points to the importance of increased cholesterol level in BMD reduction.

The association between statins and bone metabolism could be explained by mevalonate synthesis inhibition. This inhibition stops the synthesis of intermediary lipid precursors which osteoclasts use for modification and activation of intracellular proteins (3,19). This mechanism connects the cholesterol level and BMD. In our study, a weak inverse connection between BMD change and cholesterol level as well as between BMD and LDL-C in patients on statin therapy after one year of follow-up was shown. The weak association between Hol and LDL-C with BMD indicates that statins have important spectrum of pleotropic effects on bone metabolism. Anti-inflammatory effects of statins could be one of the mechanisms involved in bone protection, having in mind the importance of inflammation in pathogenesis of osteoporosis (20). This is of special importance in postmenopausal period which is followed by increase in proinflammatory cytokines (21).

Having in mind the obtained study results, we can conclude that simvastatin therapy showed a positive trend and increase in bone mineral density in hypercholesterolemic postmenopausal women after twelve months of follow-up, however, without statistically significant difference.

References

1. La Rosa JC, He J, Vupputuri S. Effect of statins on risk of coronary disease: A meta-analysis of randomized controlled trials. *JAMA* 1999; 282: 2340-6.
2. Heart Protection Study Collaborative Group: MRC /BHF Heart Protection Study of cholesterol lowering with Simvastatin in 20536 high risk individuals: A randomised placebo-controlled trial. *Lancet* 2002; 360:7-22.
3. Van Beek E, Lowik C, van der Pluijm G, et al. The role of geranylgeranylation in bone resorption and its suppression by bisphosphonates in fetal bone explants in vitro: A clue to the mechanism of action of nitrogen-containing bisphosphonates. *J Bone Miner Res* 1999;14:722-9.
4. Mundy G, Garrett R, Harris S, et al. Stimulation of bone formation in vitro and in rodents by statins. *Science* 1999;286:1946-9.
5. Reid IR, Tonkin A., Cannon CP. Comparison of the effects of pravastatin and atorvastatin on fracture incidence in the PROVE IT-TIMI 22 trial-Secondary analysis of a randomized controlled trial. *Bone* 2005;37:190 - 191.
6. Wang PS, Solomon DH, Mogun H, et al. HMG-CoA reductase inhibitors and the risk of hip fractures in elderly patients. *JAMA* 2000;283:3211-6.
7. Chan KA, Andrade SE, Boles M, et al. Inhibitors of hydroxymethylglutaryl-coenzyme A reductase and

- risk of fracture among older women. *Lancet* 2000; 355:2185-8.
8. Pasco JA, Kotowicz MA, Henry MJ, et al. Statin use, bone mineral density, and fracture risk: Geelong Osteoporosis Study. *Arch Intern Med* 2002;162: 537-540.
 9. van Staa TP, Wegman S, de Vries F, et al. Use of statins and risk of fractures. *JAMA* 2001; 285: 1850-5.
 10. Reid IR, Hague W, Emberson J, et al. Effect of pravastatin on frequency of fracture in the LIPID study: Secondary analysis of a randomised controlled trial. Long-term Intervention with Pravastatin in Ischaemic Disease. *Lancet* 2001;357:509-12.
 11. Pedersen TR, Kjekshus J. Statin drugs and the risk of fracture. 4S Study Group. *JAMA* 2000;284: 1921-2.
 12. Kanis JA, Johnell O, Oden A, et al. Risk of hip fracture according to the World Health Organization criteria for osteopenia and osteoporosis. *Bone* 2000;27:585-90.
 13. Adami S, Braga V, Gatti D. Association between bone mineral density and serum lipids in men. *JAMA* 2001;286:791-2.
 14. Chung YS, Lee MD, Lee SK, et al. HMG-CoA reductase inhibitors increase BMD in type 2 diabetes mellitus patients. *J Clin Endocrinol Metab* 2000; 85:1137-42.
 15. Edwards CJ, Hart DJ, Spector TD. Oral statins and increased bone-mineral density in postmenopausal women. *Lancet* 2000;355:2218-19.
 16. Watanabe S, Fukumoto S, Takeuchi Y, et al. Effects of 1-year treatment with Fluvastatin or Pravastatin on bone. *Am J Med* 2001;110:584-7.
 17. Montagnani A, Gonnelli S, Cepollaro C, et al. Effect of Simvastatin treatment on bone mineral density and bone turnover in hypercholesterolemic postmenopausal women: A 1-year longitudinal study. *Bone* 2003; 32:427-33.
 18. Rejnmark L, Buus NH, Vestergaard P, et al. Statins decrease bone turnover in postmenopausal women: A cross sectional study. *Eur J Clin Invest* 2002; 32:581-9.
 19. Hatzigeorgiou CC, Jackson JL. Hydroxymethyl-glutaryl-coenzyme A reductase inhibitors and osteoporosis: a meta-analysis. *Osteoporosis Int* 2005;16: 990-8.
 20. Lane NE, Rehman Q. Osteoporosis in the rheumatic disease patient. *Lupus* 2002;11:675-9.
 21. Satyawati BJ, Girish KJ. Statins and osteoporosis. *JPP* 2006, 58:3-18.

EFEKTI TERAPIJE SIMVASTATINOM NA MINERALNU GUSTINU KOSTI KOD ŽENA SA HIPERHOLESTEROLEMIJOM U POSTMENOPAUALNOM PERIODU

Todorka Savić¹, Dimitrije Janković², Irena Janković³, Vladmila Bojanić⁴, Boris Đinđić⁴,
Nataša Miladinović-Tasić⁵, Ivan Tasić¹

¹Institut za lečenje i rehabilitaciju „Niška Banja“, Srbija

²Swiss Pharm Srbija

³Klinika za plastičnu hirurgiju, Klinički centar Niš, Srbija

⁴Institut za patofiziologiju, Medicinski fakultet u Nišu, Srbija

⁵Zavod za zaštitu zdravlja Niš, Srbija

Sažetak

Cilj našeg ispitivanja bio je procena efekata terapije simvastatinom na mineralnu gustinu kosti kod žena u postmenopauzi sa hiperholesterolemijom.

U ispitivanje su uključene pedeset i tri žene sa hiperholesterolemijom u postmenopausalnom periodu, prosečne starosti 62,92+6,08 godina, koje su lečene u Institutu za lečenje i rehabilitaciju reumatičnih i kardiovaskularnih bolesti Niška Banja. Ispitanice su podeljene u dve grupe: sa vrednostima ukupnog holesterola (Hol) $\geq 7,8$ mmol/l sačinjavale su grupu I (n=32), a ispitanice sa vrednostima ukupnog holesterola manjim od 7,8 mmol/l (n=21) sačinjavale su kontrolnu grupu II. Ispitanice grupe I lečene su, pored higijensko dijetetskog režima, sa 20 mg simvastatina dnevno. Određivani su trigliceridi (Tg), ukupni holesterol (Hol), LDL holesterol (LDL-C) i holesterol u lipoproteinu velike gustine (HDL-C). Mineralna gustina kosti određivana je na lumbalnoj kičmi (L1-L4), pomoću denzitometra marke "LUNAR DPX", a rezultat je izražavan kao apsolutna vrednost (g/cm²) i T skor. Biohemijski parametri i mineralna gustina kosti (MGK) određivani su na početku i nakon 12 meseci ispitivanja.

Nakon 12 meseci primene simvastatina u dozi od 20mg/dan u grupi I došlo je do značajnog smanjenja nivoa Hol (-12,68%; $p < 0,01$) i LDL-C (-17,99%; $p < 0,005$) i do neznčajnih promena nivoa Tg (6,36%) i HDL-C (2,22%). U grupi II je nakon 12 meseci primene hipolipemijske dijeta došlo do nesigifikantnih promena lipidnih parametara. U grupi I, MGK pokazuje porast od 2,81% (NS), dok u grupi II dolazi do njenog smanjenja za -3,45% (NS). Kod ispitanica koje su bile na statinskoj terapiji utvrđena je negativna korelacija između promena MGK i LDL-C ($r = -0,08$; $p = 0,686$; $r = 0,166$, $p = 0,626$), kao i između promena MGK i Hol ($r = -0,144$; $p = 0,476$; $r = -0,125$; $p = 0,715$) tokom godinu dana ispitivanja. Ovi rezultati ne pokazuju statistički značaj i nisu prikazani grafički.

Terapija simvastatinom dovodi do trenda porasta mineralne gustine kostiju kod žena sa hiperholesterolemijom u postmenopauznom periodu života nakon 12 meseci terapije, ali bez statistički značajnog porasta.

Ključne reči: mineralna gustina kostiju, statinska terapija, žene u postmenopauzi