CENTRAL CORNEAL THICKNESS MEASURED BY THE OCULYZER, BIOGRAPH, AND ULTRASOUND PACHYMETRY

Marko Zlatanović¹, Maja Živković^{1,2}, Aleksandra Hristov³, Vesna Stojković³, Saša Novak¹, Nevena Zlatanović², Mladen Brzaković³

¹Ophthalmology Clinic, Clinical Center, Niš, Serbia ²University of Niš, Faculty of Medicine, Niš, Serbia ³Specialized Eye Hospital "Maja Clinic" Niš, Serbia

Contact: Aleksandra Hristov Djordja Andrejevića Kuna 24, Niš E-mail: alexandramiljkovic@gmail.com

Central corneal thickness (CCT) is an important parameter in selecting patients before any corneal refractive procedures and in postoperative follow-up, in monitoring the patients with corneal disorders such as keratoconus and Fuchs' dystrophy, as well as in adjusting the intraocular pressure according to central corneal thickness.

The aim of the paper was to compare the results of CCT measurements obtained by different measurement instruments: Oculyzer, BioGraph and ultrasound pachymetry.

CCT was measured in 64 eyes of 32 patients. The measurements were performed in a specialized Eye Hospital "Maja Clinic" in Niš, using the following instruments: WaveLight Allegro Oculyzer, WaveLight Allegro Biograph, ultrasound pachymeter DGH Pachette 3. The mean values of central corneal thickness obtained from different devices were compared.

The mean values of CCT (± standard deviation) obtained with Oculyzer were 552.94 μ m ± 22.88 μ m, 556.56 ± 25.32 μ m obtained with BioGraph, and 559.46 ± 26.0 μ m measured by ultrasound pachymeter. There were no statistically significant differences among different measurement devices.

The CCT measurements with the Oculyzer, BioGraph, and ultrasound pachymetry do not show statistically significant differences, so the results of CCT measurements obtained by using any of these devices may be considered valid.

Acta Medica Medianae 2019;58(2):33-37.

Key words: corneal thickness, pachymetry, measurement devices