

THE IMPORTANCE OF MAGNETIC RESONANCE IMAGING IN THE DIAGNOSIS OF TEMPOROMANDIBULAR DISORDERS

Vladimir Rakić¹, Vladimir Antić², Milorad Antić³

The addition of new imaging modalities, specifically magnetic resonance imaging, has and will continue to increase the amount of diagnostic information available to the clinician. Imaging is the only method of obtaining visual information on the status of the joint tissues short of arthroscopy or open joint surgery. Magnetic resonance imaging has capacity to contrast soft tissue types makes it an ideal tool for investigating internal derangements. Its ability to image in multiple planes is well suited for examining the three-dimensional nature of internal derangements of temporomandibular joints. A major bonus is its freedom from ionizing radiation and other known health hazards making it a low-risk method for research and clinical investigation. A major disadvantage is its cost and associated limited availability. Magnetic resonance imaging examinations of the temporomandibular joints are part of the current standard of care in the evaluation of temporomandibular disorders, providing the clinician anatomic information that can guide treatment decisions.

Acta Medica Medianae 2019;58(4):xx-xx.

Key words: magnetic resonance imaging, temporomandibular joint, temporomandibular disorders

¹Department for Radiology, Clinical center Niš, Serbia

²University of Niš, Faculty of Sport and Physical Education, Niš, Serbia

³University of Niš, Medical faculty, Institute of Anatomy, Niš, Serbia

Contact: Vladimir Rakić
Blvd dr Zoran Djindjić 48, 18 000 Niš, Serbia
E-mail: vladimir_vlada@yahoo.com

Introduction

The radiologic investigation of the temporomandibular joints may bring to light pertinent information. However, only after relating these findings to the clinical symptoms will the diagnostic significance become apparent. Since some osseous changes may not be detectable in the radiologic examination, the final diagnosis becomes a clinical procedure. The addition of new imaging modalities, specifically magnetic resonance imaging (MRI) has and will continue to increase the amount of diagnostic information available to the clinician (1). Imaging is the only method of obtaining visual information on the status of the joint tissues short of arthroscopy or open joint surgery. Its primary

purpose is to provide information to assist the diagnosis and treatment planning process. Despite temporomandibular joint imaging's long history of research and clinical application, the quality of information gleaned from imaging is often less than desired (2). The small size of the TMJ, the widely varying fossa and condylar morphology and the surrounding dense osseous structures make clear and undistorted imaging of the joint hard tissue technically difficult (3).

TMJ anatomy and function

Major components of the temporomandibular joint include the mandibular condyle, the articular disc, the glenoid fossa, and the articular eminence of the temporal bone. Unlike most joints, the articulating surfaces are fibrous and not cartilaginous. The fibrocartilaginous articular disc is biconcave, dividing the joint space into superior and inferior compartments; this relationship is well seen in the presence of joint effusion (4). The anterior and posterior portions of the articular disc, which are thickened by the morphology of the disk annulus, are designated the anterior band and the posterior band, respectively, with a thinner intermediate zone in between. The disc is attached to the temporal bone and condyle posteriorly by elastic and loose connective tissue; this tissue is also known as the retrodiscal soft tissue or the bilaminar zone. The lateral pterygoid muscle, the only muscle of mastication serving to open the jaw, inserts on the

mandibular condyle inferior to the articular surface but can partially insert on the joint capsule and disc as well (5).

Magnetic resonance imaging technology

Magnetic resonance imaging technology exploits the varying proton content of different tissues. The protons in tissue fluids are polar, analogous to tiny bar magnets, with their magnetic fields or dipoles aligned in random fashion (6). When exposed to the strong magnetic field of the Magnetic resonance imaging scanner, some of the protons align parallel with direction of the external field. Radio waves of a specific frequency (similar to broadcast signals) are directed at the tissue inducing proton precession, a motion similar to toy top winding down. The magnitude of precession is proportional to the amount of radio frequency energy absorbed. When the radio frequency excitation is stopped, the protons relax to their original low energy state and, in the process, emit the absorbed energy which can be detected by receiver antennae placed over the areas of interest (7).

Magnetic resonance imaging examination of the temporomandibular joint has gained an important role in the diagnosis of internal derangement, because it allows direct visualization of the articular disk in both the open- and closed-mouth positions. Nuclear magnetic resonance was introduced 40 years ago as a research tool in chemistry and physics. The development of large superconducting magnets and high speed computers has paved the way for adaptation of the technology to clinical diagnosis. The technology as applied to medicine was renamed "magnetic resonance imaging" to avoid the stigma attached to the term "nuclear" (8). Its key advantages over other imaging techniques are elimination of ionizing radiation and the capacity to produce high resolution images in most anatomic planes (axial, sagittal, frontal and oblique), without positioning of the patient as required with direct sagittal CT scanning.

Specially small coils placed over both temporomandibular joints areas enhance the clarity of received signals and allow both joints to be imaged in a single exam sequence (9). The strength of the emitted signals is proportional to the amount of protons in the tissue. Signals are location coded as a result of strength gradation in the primary magnetic field (10). This allows the computer to assign an intensity and location values to the emitted signals which are then manipulated by the computer into cross-sectional images.

Magnetic resonance imaging is most commonly applied to the diagnosis of internal derangements (11). However, it has potential to diagnose hard tissue lesions. Some authors reported

that number of bony abnormalities were noted on coronal views that were not appreciated on sagittal views nor in some case, on tomography. Magnetic resonance imaging has been reported to be 95% accurate in assessment of disk position and form and 93% accurate in assessment of osseous changes (12). However, several authors have noted a lack of correlation between magnetic resonance imaging findings of disk displacement and the extent of pain and dysfunction of the temporomandibular joint in patients with painful limitation of mandibular opening (13,14). Moreover, disk displacement was found in a substantial number of asymptomatic volunteers. For example, Ahmad and colleagues reported a 21% prevalence of internal derangement on magnetic resonance imaging evaluation of 57 asymptomatic people (15).

The clinical significance of imaging findings of internal derangement is controversial (16). The prevalence of displacement of the temporomandibular joint disk among asymptomatic volunteers was previously reported as nearly 33% and the prevalence of normal articular disk in symptomatic joints was reported to be 16%-23% (17). Moreover, arthroscopy and magnetic resonance imaging have shown that temporomandibular joints with anteriorly displaced disks have the capacity to form remodelled retrodiscal tissue that resembles cartilage (i.e., pseudo-disk formation) (18). Furthermore, the retrodiscal tissues have adaptive capacity and often respond appropriately to the functional loads placed on the tissues (19,20).

Conclusion

Magnetic resonance imaging has capacity to contrast soft tissue types makes it an ideal tool for investigating internal derangements. Its ability to image in multiple planes is well suited for examining the three-dimensional nature of internal derangements of temporomandibular joints. A major bonus is its freedom from ionizing radiation and other known health hazards making it a low-risk method for research and clinical investigation. A major disadvantage is its cost and associated limited availability. Magnetic resonance imaging examinations of the temporomandibular joints are part of the current standard of care in the evaluation of temporomandibular disorders, providing the clinician anatomic information that can guide treatment decisions. This article has reviewed some of the key findings and imaging appearances of the degenerated temporomandibular joint. Further research will continue to enhance our understanding of the potential contributions of contrast-enhanced studies and dynamic imaging.

References

1. Sidebottom AJ. Current thinking in temporomandibular joint management. *Br J Oral Maxillofac Surg*. 2009;47(2):91-4.
2. Emshoff R, Brandlmaier I, Gerhard S, Strobl H, Bertram S, Rudisch A. Magnetic resonance imaging predictors of temporomandibular joint pain. *J Am Dent Assoc*. 2003;134(6):705-14.
3. Schmitter M, Kress B, Ludwig C, et al. Temporomandibular joint disk position assessed at coronal MR imaging in asymptomatic volunteers. *Radiology*. 2005;236:559-64.
4. Sano T, Westesson PL, Larheim TA, Takagi R. The association of temporomandibular joint pain with abnormal bone marrow in the mandibular condyle. *J Oral Maxillofac Surg*. 2000;58:254-57.
5. Taskaya-Yilmaz N, Ceylan G, Incesu L, Muglali M. A possible etiology of the internal derangement of the temporomandibular joint based on the MRI observations of the lateral pterygoid muscle. *Surg Radiol Anat*. 2005;27(1):19-24.
6. Wang EY, Mulholland TP, Pramanik BK, et al. Dynamic sagittal half-Fourier acquired single-shot turbo spin-echo MR imaging of the temporomandibular joint: Initial experience and comparison with sagittal oblique proton-attenuation images. *Am J Neuroradiol*. 2007;28:1126-32.
7. Scrivani SJ, Keith DA, Kaban LB. Temporomandibular disorders. *The New England Journal of Medicine*. 2008;359:2693-05.
8. Kubein-Meesenburg D, Nägerla H, Fialka-Fricke J, Hahn W, Weber S, Hönig J, Hansen C, Fanghänel J, Thieme KM, Ihlow D. Functional states of mandibular movements and synovial pumps of the temporomandibular joint. Is it possible to provide a biomechanically correct replacement for the TMJ. *Ann Anat*. 2012;194(2):200-7.
9. Macfarlane TV, Blinkhorn AS, Davies RM, Kincey J, Worthington HV. Oro-facial pain in the community: prevalence and associated impact. *Community Dent Oral Epidemiol*. 2002;30:52-60.
10. Macfarlane TV, Kenealy P, Kingdon HA, Mohlin B, Pilley JR, Mwangi CW, Hunter L, Richmond S, Shaw WC. Orofacial pain in young adults and associated childhood and adulthood factors: results of the population study, Wales, United Kingdom. *Community Dent Oral Epidemiol*. 2009;37:438-50.
11. Manfredini D, Guarda-Nardini L, Winocur E, et al. Research diagnostic criteria for temporomandibular disorders: a systematic review of axis I epidemiologic findings. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:453-62
12. Li C, Zhang Y, Lv J, et al. Inferior or double joint spaces injection versus superior joint space injection for temporomandibular disorders: a systematic review and meta-analysis. *J Oral Maxillofac Surg* 2012;70(1):37-44.
13. Westesson PL, Brooks SL. Temporomandibular joint: relationship between MR evidence of effusion and the presence of pain and disk displacement. *Am J Roentgenol*. 1992;159:559-63.
14. Peck CC, Goulet JP, Lobbezoo F et al. Expanding the taxonomy of the diagnostic criteria for temporomandibular disorders. *J Oral Rehabil*. 2014;41:2-23.
15. Ahmad M, Schiffman EL. Temporomandibular Joint Disorders and Orofacial pain. *Oral Radiology: Dent Clin N Am*. 2016;60: 105-24.
16. Yura S, Harada S, Kobayashi K. Diagnostic Accuracy on Magnetic Resonance Imaging for the Diagnosis of Osteoarthritis of the Temporomandibular Joint. *J Clin Diagn Res*. 2015;9:95-7.
17. Murakami K. Rationale of arthroscopic surgery of the temporomandibular joint. *J of Oral Biol and Craniofac Res*. 2013;3 :126-34.
18. Aiken A, Bouloux G, Hudgins P. MR imaging of the temporomandibular joint. *Magn Reson Imaging Clin N Am*. 2012;20:397-412.
19. Sava A, Scutariu M. Functional anatomy of the temporo-mandibular joint (II) *Rev Med Chir Soc Med Nat Iasi*. 2012;116:1213-17
20. Imanimoghaddam M, Madani AS, Hashemi EM. The evaluation of lateral pterygoid muscle pathologic changes and insertion patterns in temporomandibular joints with or without disc displacement using magnetic resonance imaging. *Int J Oral Maxillofac Surg*. 2013;42:1116-20.

ZNAČAJ MAGNETNE REZONANCE TEMPOROMANDIBULARNOG ZGLOBA U DIJAGNOSTICI TEMPOROMANDIBULARNIH POREMEĆAJA

Vladimir Rakić¹, Vladimir Antić², Milorad Antić³

¹Odeljenje radiologije, Klinički centar Niš, Srbija

²Univerzitet u Nišu, Fakultet sporta i fizičkog vaspitanja, Niš, Srbija

³Univerzitet u Nišu, Medicinski fakultet, Institut za anatomiju, Srbija

Kontakt: Vladimir Rakić
Bulevar dr Zorana Đinđića 48, 18000 Niš, Srbija
E-mail: vladimir_vlada@yahoo.com

Uvođenje novih načina rendgenskih snimanja, posebno magnetne rezonance, pruža sve veći broj dijagnostičkih informacija koje su dostupne kliničarima. Magnetna rezonanca je jedan od načina dobijanja vizuelnih informacija o stanju zglobnog tkiva bez artroskopije ili otvorene operacije zgloba. Magnetna rezonanca ima sposobnost da pomoću kontrastnih supstanci određuje tipove mekih tkiva, što ga čini idealnim alatom za istraživanje unutrašnjih poremećaja. Njena sposobnost da daje slike u više ravni je pogodna za ispitivanje trodimenzionalne prirode unutrašnjih poremećaja temporomandibularnih zglobova. Glavna prednost je odsustvo jonizujućeg zračenja i drugih agenasa opasnih po zdravlje, što ga čini metodom niskog rizika za istraživanje i kliničko ispitivanje. Glavni nedostatak je njegova cena i ograničena dostupnost. Ispitivanja temporomandibularnih zglobova magnetnom rezonancom su deo standarda u proceni temporomandibularnih poremećaja, pružajući kliničarima anatomske informacije koje mogu uticati na donošenje konačne odluke o lečenju.

Acta Medica Medianae 2019;58(4):xx-xx.

Ključne reči: snimanje magnetnom rezonancom, temporomandibularni zglob, temporomandibularni poremećaji