

BERRIES AS A NATURAL SOURCE OF BONE SUPPORT: THE INVOLVEMENT OF ANTHOCYANINS IN THE MOLECULAR MECHANISMS OF THE HEALING AND REGENERATION PROCESSES

Jelena Najdanović^{1,2}, Stevo Najman^{1,2}, Jelena Živković^{1,2},
Bojana Miladinović³

Bone tissue has a remarkable self-regenerating ability which, in case of injury, enables return to a completely functional, pre-injury state. However, ageing, disease, obesity, traumas, multiple fractures, infections and tumor removal cause large bone defects that cannot be healed spontaneously. To achieve successful bone healing and regeneration, plenty of approaches, including the application of autografts, allografts and bone tissue engineering (BTE), have been developed. One of the approaches is based on the findings that bone loss in humans and many animals during aging is partially caused by accumulation of reactive oxygen species (ROS). Due to the spectrum of biological activities, including antioxidative, essential polyphenolic components—anthocyanins (ACNs), are a part of a significant research area regarding means and methods for bone healing and regeneration. Berries are especially rich in ACNs. Based on *in vitro* and *in vivo* studies regarding molecular mechanisms involved in bone healing and regeneration supported with berries' ACNs and on observational research in human populations, it has been found that berries' ACNs enhance osteoblastogenesis, suppress osteoclastogenesis and have osteoimmunological activity. Therefore, berries' ACNs should be considered as naturally widespread therapeutics for bone support. Nevertheless, before implementation of berries as a natural source of bone support, there are some issues left to resolve: clarification of molecular mechanisms of ACNs action in bone metabolism, identification of effective doses of particular ACNs for bone regeneration therapies and performing clinical studies for determination of therapeutic efficacy of different types and concentrations of ACNs.

Acta Medica Medianae 2024;63(4):104–113.

Key words: bone healing, bone regeneration, berries, anthocyanins, molecular mechanisms

¹University of Niš, Faculty of Medicine, Department of Biology and Human Genetics, Niš, Serbia

²University of Niš, Faculty of Medicine, Department of Cell and Tissue Engineering, Scientific Research Center for Biomedicine, Niš, Serbia

³University of Niš, Faculty of Medicine, Department of Pharmacy, Niš, Serbia

Contact: Jelena Najdanović
81 Dr. Zorana Djindjića Blvd., 18000 Niš, Serbia
E-mail: jelena.najdanovic@medfak.ni.ac.rs

adulthood to a certain degree which, in case of injury, enables the return to a completely functional, pre-injury state (2). However, a high rate of bone tissue degradation and decreased production of new bone lead to bone loss (1). Bone mass declines and bone microarchitecture weakens with ageing so that, in advanced age, the bone resorption rate surpasses the bone formation rate (3). Besides ageing, other systemic factors, such as disease and/or obesity, also have an influence on bone healing and regeneration (4). Moreover, traumas, multiple fractures, infections and tumor removal cause large defects that cannot be healed spontaneously (5).

Bone healing and bone regeneration approaches

To achieve successful bone healing and regeneration, plenty of approaches have been developed. The gold standard in the treatment of posttraumatic conditions—fractures, delayed unions and nonunions, is the application of autografts—the grafts constructed out of the

Introduction

Bone-forming osteoblasts (OBs) and bone-resorbing osteoclasts (OCs) control normal bone metabolism (1). Remarkable bone tissue self-regenerating ability is maintained throughout

patient's own bone (6). Autografts' structure is similar to the original bone, so bone growth and regeneration are enabled (7), but taking autologous bone is usually associated with health risks to the patient and the source of the tissue is limited (8). Processed bone allografts are good alternatives because of their higher availability and the lack of donor-site complications in the recipient (6). However, the limitation of allografts is the induction of immune reactions which is followed by graft rejection, the possibility of infection and the chance of disease transmission (9). In the field of bone tissue engineering (BTE), great efforts have been made to avoid complications caused by bone grafts. The idea of BTE is to mimic the structure of a natural bone and construct an implant based on a biological triad—biomaterial as a scaffold, regulatory molecules and cells (10). In light of that, implants based on combinations of different biomaterials, various regulatory molecules and different types of cells were constructed. The promising results in BTE include the application of calcium phosphate-based biomaterials as a scaffold due to their similarity with the natural bone structure (11, 12, 13). Also, the inclusion of natural sources of growth factors, such as platelet-rich plasma (PRP) (14) and blood (15) into the implants' structure deserves special attention in BTE. Biomaterials and regulatory molecules can be combined with macrophages (13, 16) or stem cells, including adipose tissue-derived mesenchymal stem cells (ADSCs). ADSCs can be applied either freshly isolated (17), *in vitro* expanded before implantation (18) or *in vitro* differentiated into various types of cells, including OBs and endothelial cells, before implantation (11, 12, 18).

Another approach for bone healing and regeneration is based on findings that bone loss in humans and many animals during ageing is partially caused by the accumulation of reactive oxygen species (ROS) (19). Cells normally prevent the excessive presence of ROS thanks to their antioxidant defense system which includes vitamins, enzymes and other substances (20). Excessive ROS accumulation leads to DNA damage, lipid peroxidation and oxidation of amino acids with consequent changes in the cells' structures and functions (21). In bone tissue, oxidative stress leads to OBs and osteocyte apoptosis (22) and stimulates osteoclastogenesis (19). During osteoclastogenesis, the transcription factor nuclear factor-kappa B (NF- κ B) signaling pathway plays a crucial role (1). The existence of such an association between oxidative stress and bone mass loss during ageing points out to the need to improve an individual's antioxidant defense. Standard pharmacological agents that improve bone mass and reduce fractures may show side effects and/or poor efficacy in bone healing and regeneration (23).

Folk medicine has various approaches to bone healing, and some of them are a source of valuable facts for creating more effective modern

bone treatments. For example, water extracts of flavonoid-rich plants, such as *Labisia pumila* (Blume) Fern.-Vill. (Myrsinaceae) and *Piper sarmentosum* Roxb. (Piperaceae), are used in folk medicine by Malay women since these plants maintain the estrogen level at the post-menopausal stage and consequently increase bone formation and reduce bone resorption (24). Root barks and stem barks rich in flavonoids are traditionally used for bone fracture healing by the indigenous people of Eastern Ghat (25). Several flavonoid-rich plants, including *Pholidota articulata* Lindl. and *Coelogyne cristata* Lindl. (Orchidaceae), are part of folk tradition in India for the treatment of bone-related disorders and fractured bones (26).

In recent decades, there has been a noticeable trend in the application of a suitable diet for the additional treatment of certain bone conditions and diseases. All-natural nutritional therapies are expected to be safer therapeutic options for bone loss and restoration of normal bone metabolism. A plethora of preclinical and clinical researches indicate that fruits and vegetables-rich diets could help bone fracture healing (27). Due to the spectrum of observed biological activities, flavonoid substances, especially anthocyanins (ACNs) as bioactive components of natural origin, are part of a significant research area regarding means and methods for bone healing and regeneration.

Anthocyanins: classifications, biological activities and health effects

With around 8000 polyphenols among which nearly 500 are bioactive, plants represent an abundant source of antioxidants (28). Berries are especially rich in polyphenols. Due to the health benefits associated with polyphenols, the intake of berries and the possibility of their application in medicine have gained much interest within scientific circles (29). Phenolic compounds have a spectrum of biological activities including anticancer, antidiabetic, anti-inflammatory, and anti-platelet, and represent one of the most powerful natural antioxidants (30).

Polyphenols are categorized into four groups: phenolic acids, flavonoids, stilbenes, and lignans (31). It has been found that flavonoids can promote bone formation and stimulate osteogenic differentiation of mesenchymal stem cells (MSCs) (32). The idea for the application of flavonoids for bone healing and regeneration in clinical medicine comes from the experience of using flavonoid-rich compounds in folk medicine worldwide (33).

Flavonoids can be classified into flavones, flavonols, isoflavones, flavanones, flavanes, chalcone, isoflavanes and ACNs (34). ACNs are essential polyphenolic water-soluble plant pigments which consist of an anthocyanidin (aglycone) bound to sugar fraction (35). According to the position and number of hydroxyl and methoxy groups, more than 635 ACNs have been

identified (1). Cyanidins are the most abundant group of ACNs found in food, afterwards delphinidins, pelargonidins, peonidins, malvidins, and petunidins (36). These plant pigments are responsible for the wide range of colors (from dark red to blue) visible to the human eye. ACNs are indicators of the ripeness as well as the quality of the fruit (37, 38). The berry skin contains the highest amount of ACNs, although they are also present in the pulp (39, 40). Their production is affected by various environmental conditions, such as light, temperature, presence of minerals, climate and many other factors.

In vitro and *in vivo* studies indicate that ACNs have anti-inflammatory and antioxidative properties (41) that prevent or delay the onset of chronic diseases involving oxidative stress and inflammation (29, 42). ACNs stop pro-inflammatory mediators by blocking their production or activity, which is the foundation of their anti-inflammatory potential (43). Some types of ACNs possess ROS scavenging properties thus preventing DNA damage (44).

Among berries, blackcurrant (*Ribes nigrum* L.) stands out as a rich source of ACNs (45), with 250 mg of ACNs per 100 g of fresh fruit (46) and ACNs' concentrations that are up to four-fold higher in comparison with other similar fruits (47). Blackcurrant contains delphinidin-3-O-rutinoside (del-3-rut), cyanidin-3-O-rutinoside (cya-3-rut), delphinidin-3-O-glucoside (del-3-glc) and cyanidin-3-O-glucoside (cya-3-glc), contributing to approximately 98% of total ACNs (48). Delphinidins contribute to approximately 74% of total ACNs in blackcurrant. They are of particular importance for the prevention of bone resorption (49) due to the higher free-radical scavenging capacity in comparison with all other major ACNs (50).

A review of the literature revealed other berries valuable as a beneficial source of ACNs. Blueberries and grapes are abundant in delphinidin, cyanidin, petunidin and malvidin glycosides, making up over 90% of the total ACN content (51, 52). Malvidins account for 16% of the total blueberry ACNs (48), whereas strawberries' attractive color and health benefits are derived from pelargonidin-3-O-glucoside as the major ACN of this popular red berry (38, 53). Cranberries' dominant ACNs are peonidin glycosides (54), whilst cyanidin glycosides are most represented in red currants, blackberries and raspberries (48, 55, 56). The role of these ACNs in bone healing and regeneration will be thoroughly elucidated in this paper.

Anthocyanins in bone healing and regeneration: mechanisms of action

Discovering the possibilities for bone healing and regeneration supported with berries' ACNs opens the door to explore the potential use of ACNs in therapeutic interventions for people who suffer from bone degeneration related to

inflammation, menopause or ageing (57). Potential benefits of ACNs are especially important for those populations that are more and more prone to osteoporosis (57), the most common bone disease characterized by low bone mineral density (BMD) and bone matrix fragility that predisposes patients to increased risk of fractures (28, 58). Osteoporosis incidence is influenced by various factors including ageing, insufficient estrogen levels, increased oxidative stress, chronic inflammation, and genetics (58).

Human (59) and animal (60) studies have shown that a strong positive correlation exists between excessive ROS and bone loss. ROS directly contribute to bone degradation by osteoclast (OC)-generated superoxide or can cause an increase in OCs' differentiation and function (60). At the same time, excessive ROS inhibit osteogenic differentiation through extracellular signal-regulated kinases (ERK) and ERK-dependent NF- κ B signaling pathways (22). Also, an important factor in the oxidant-antioxidant balance is the capacity of OBs to produce antioxidants in response to ROS (61).

It has been reported that ACNs can enhance osteoblastogenesis, suppress osteoclastogenesis and also have osteoimmunological activity (28). The main modes of ACNs' action on osteogenesis are achieved by involving in the molecular mechanisms of bone morphogenetic protein (BMP-2), WNT- β catenin and fibroblast growth factor (FGF). Del-3-rut and cya-3-glc activate the FGF pathway thus accomplishing the influence on OBs' differentiation. Most of the ACNs that can promote osteogenesis also up-regulate the expression of genes for transcription factors Sox9, Runx2, and Osterix (Osx) and genes for type 1 collagen (Col1), osteopontin (OPN), osteocalcin (OCN), and alkaline phosphatase (ALP) (1). The effects of different ACNs on osteoclastogenesis are mainly achieved by involvement in the molecular mechanisms of some pathways such as c-Fos, NF- κ B, JNK, Ca²⁺ and ROS (1). In addition, three subfamilies of mitogen-activated protein kinases (MAPKs) are important in RANK signal-mediated OC generation (62), while the nuclear factor of activated T-cells 1 (NFATc1) represents a major pathway that regulates osteoclastogenesis.

Bone healing and regeneration supported with ACNs and berries compounds: *in vitro* studies

The involvement of ACNs and berries compounds in bone healing and regeneration was examined by using various *in vitro* models. One type of applied *in vitro* models mimic the conditions in an organism that are the consequence of estrogen deficiency and/or microdamage. In one of such models, oxidative stress was induced in the human OB-like cell line SaOS-2 by an intracellular depletion of glutathione (GSH), in the period before the beginning of osteogenic differentiation and during the early

mineralization process (63). In GSH-depleted SaOS-2 cells, blueberry juice (BJ) rich in ACNs prevented inhibition of osteogenic differentiation and mineralization process caused by oxidative stress. BJ also modulated signals which up-regulate the expression and activity of osteogenic factors. Likewise, increased expression of sirtuin type 1 deacetylase, an enzyme that regulates osteogenic differentiation of tendons and MSCs and represents a positive Runx2 regulator (64), is probably related to the osteogenic action of BJ (63).

Inhibitory effects of delphinidin, cyanidin and peonidin on osteoclastogenesis were examined and compared—delphinidin suppressed *in vitro* OC formation, while cyanidin and peonidin did not showed such strong impact on osteoclastogenesis (58). On the other hand, Ostos Mendoza et al. (65) revealed that, peonidin-3-O-glucoside applied in low concentrations improved OBs' viability and reduced apoptosis in serum-starved human OBs. This treatment favored the cell growth and OBs differentiation as well as the alteration in the expression of proinflammatory interleukins and downregulation of RANKL expression, which suggests the possible use of peonidin as therapeutic in bone diseases.

In vitro osteogenic effects of delphinidin glycoside-enriched maqui berry extract (MBE) were also examined (66). Up-regulated bone-related gene expression for proteins such as BMP-2, OSX, and OCN indicated that MBE stimulated osteogenic differentiation of MC3T3-E1 cells. The research conducted on RAW264.7 cell line showed that petunidin (> 5 µg/ml) significantly suppressed OCs' differentiation and down-regulated expression of genes for c-Fos, NFATc1, matrix metalloproteinase 9 and cathepsin K (67).

The anti-inflammatory effects of ACNs contained in blueberry, blackberry and blackcurrant were compared, and the relationship between their antioxidant capacity and anti-inflammatory effect in macrophages was determined (48). These berries achieved anti-inflammatory effects in macrophages, at least partially, due to inhibition of nuclear translocation of NF-κB independent of the nuclear factor E2-related factor 2 (NRF2)-mediated pathways.

Multiple effects of cya-3-glc on OCs are well-known, but the mechanisms of its impact on OBs are not yet completely clarified. Therefore, the effects of cya-3-glc on proliferation and differentiation of the hip joint-derived OBs taken from osteoporotic patients and on mice OB cell line MC3T3-E1 were examined (68). The ability of OBs to mineralize after cya-3-glc treatment as well as the role of ERK signaling pathway in cya-3-glc regulation of OBs were also evaluated. ERK, a crucial member of MAPK cascades, positively regulates OB differentiation and bone formation (69). Cya-3-glc enhanced OBs' proliferation rate and OBs' mineralization points, up-regulated OCN gene and protein expression, and increased the level of ERK phosphorylation (68), which proves

that ERK pathway is involved in cya-3-glc regulation of osteogenic differentiation and indicates that OBs can be targets for prevention and treatment of osteoporosis.

Del-3-rut protects MC3T3-E1 from oxidative damage and promotes osteogenic differentiation of this cell line via PI3K/AKT pathway (1), which means that del-3-rut could be used as dietary supplement for the prevention of OBs' dysfunction in age-related osteoporosis (70). Also, malvidins are considered to be responsible for bone formation by inducing significantly higher calcium deposits in MSCs (71).

Bone healing and regeneration supported with ACNs and berries compounds: *in vivo* studies

Numerous *in vivo* studies reveal a valuable relationship between ACNs and bone health. Bone-protective roles of phenolic and flavonoid ingredients derived from dried plum have been shown in rat osteoporosis models (72). The fact that dried plum and blueberry have several identical phenolic and flavonoid ingredients was used to perform research on an ovariectomized (OVX) rat model of postmenopausal osteoporosis (42). It was hypothesized that blueberry-derived phenolic compounds can prevent bone loss in ovarian hormone deficiency (42). OCN, Col1 and bone-specific ALP were chosen as markers of bone formation, and tartrate-resistant acid phosphatase (TRAP) as a bone resorption marker. Analyses at gene and protein expression levels indicated that treatment with 5% blueberry (w/w) prevented bone loss by suppression of ovariectomy-caused bone turnover. The OVX rat model was also applied to examine the effects of rabbiteye blueberry on osteoporosis (73). Rabbiteye blueberries effectively inhibited bone resorption, bone loss, and reduction of bone strength of OVX rats. In another study, blackcurrant extract supplementation reduced trabecular and cortical bone loss in an OVX mice model (74), which was the same effect estimated upon supplementation of OVX mice with delphinidin glycoside-enriched MBE (66). On the other hand, bilberry extract that has 15 various ACNs did not show an impact on bone metabolism in the OVX rat model (19).

Besides the OVX model, which mimics postmenopausal estrogen loss but does not specifically mimic the effects of ageing, an age-related model of bone loss can also be used to evaluate the influence of ACNs on bone loss prevention. In this model, the influence of blackcurrant extracts on the improvement of mice bone mass was evaluated (75). Young and old female C57BL/6J mice were fed with either a standard chow diet or a chow diet enriched with 1% (w/w) blackcurrant extract for four months. Since supplementation with blackcurrant extracts improved glutathione peroxidase and catalase activity and led to an increase in trabecular bone volume, OB surface, and bone mineral content in

young mice, it was concluded that consumption of blackcurrant early in life—when a substantial amount of bone mass is still present, could prevent ageing-associated bone loss.

Besides being the direct bone remodeling mechanisms, high oxidative stress and chronic inflammation can also lead to obesity and, consequently, to bone loss. Low levels of vitamins C and E, carotenoids, superoxide dismutase, glutathione peroxidase, catalase and other plasma antioxidants and antioxidant enzymes in obesity can cause augmented bone resorption (76). Therefore, the effects of blueberry, blackberry and blackcurrant on bone health were examined on a diet-induced obesity mice model. High-fat (HF) diet-induced obese C57BL mice were fed a HF diet, with or without berry supplementation, for 12 weeks. The results confirmed that there was a negative correlation between fat and bone mass, but that consumption of berries with different ACNs' composition could affect bone turnover via mechanisms that should be clarified in the future (76).

It was discovered that pelargonidin-3-O-glucoside acts as an anti-inflammatory agent by suppressing the NF- κ B pathway in an experimental model of osteoarthritis. In this way, the inflammation and cartilage damage can be reduced as well as the progression of osteoarthritis (77). Petunidin prevented bone mass loss in a RANKL-induced osteopenic mice model (67). Cyanidin-chloride (CC) and cya-3-glc, can regulate bone homeostasis, but the literature data regarding their specific role in osteoclastogenesis are controversial. According to Cheng et al. (78), CC inhibits osteoclastogenesis, hydroxyapatite resorption, and RANKL-induced signal pathways *in vitro* and protects against OVX-induced bone loss *in vivo*. Other data indicate that, at high doses ($> 10 \mu\text{g/ml}$), cyanidins suppress osteoclastogenesis and OCs fusion, but at low doses ($< 1 \mu\text{g/ml}$) the effect is the opposite (79). Moreover, cya-3-glc improved OBs proliferation and up-regulated OCN gene and protein expression, mainly via the ERK1/2 pathway (68).

The application of flavonoids, including ACNs, in the field of BTE is becoming an increasingly attractive way to promote bone healing. Their role, in addition to protecting cells from oxidative stress, is also reflected in the promotion of proliferation and osteogenic differentiation of MSCs (80). Attempts to incorporate flavonoids into different types of biomaterials to promote bone defects' healing have proven to be more than successful. Their beneficial effect is reflected in the increase in osteogenic and angiogenic markers' expression, activation of the Wnt signaling pathway and reduction of inflammatory factors' levels (34).

The effects of anthocyanins and other flavonoids on bone healing and regeneration in population studies

The information regarding the effects of different flavonoid subclasses on bone health in humans are limited. An observational research conducted in the group of 3160 female twins revealed that total flavonoid intake was positively correlated with BMD of the hip and spine (81). Also, a strong positive correlation between consumption of fruits and BMD and bone mineral content (BMC) in boys and girls (11–14 years), young women (20–34 years), and postmenopausal women (50–70 years) has been reported (82). In addition, positive association between high fruit intake and high BMD in men and women aged 25–64 years was found (83). Another study was conducted only in women—women who consumed high amounts of fruit in childhood had higher BMD of the femoral neck compared to the women who had a medium or low intake of fruits during childhood (84).

Conclusion

Based on *in vitro* and *in vivo* studies regarding molecular mechanisms involved in bone healing and regeneration supported with berries' ACNs and on observational research in human populations, it is unequivocally clear that berries' ACNs could be used for the prevention and treatment of the certain bone conditions and diseases. Berries' ACNs are part of nature, so they should be considered as naturally widespread therapeutics for bone support. Furthermore, research into the possibility of bone healing and regeneration using substances from natural sources, such as ACNs, may contribute to the development of new, less invasive therapeutic methods. However, some discrepancies are noticed regarding the influence of ACNs on bone healing and regeneration which can be explained by the different experimental models that were chosen, different classes of ACNs that were used, or different concentrations of the same type of applied ACNs. Despite that, plenty of encouraging results speak in favor of the medical use of ACNs in bone healing and regeneration as a safer and cheaper solution for human health in comparison with standard medical therapies. Nevertheless, before the implementation of berries as a natural source of bone support, there are some issues left to resolve. First, further studies regarding clarification of molecular mechanisms of ACNs' action in bone metabolism are needed. Then, the identification of effective doses of particular ACNs for bone regeneration therapies needs to be determined. Finally, clinical studies for the determination of the therapeutic efficacy of different types and concentrations of ACNs must be performed. Resolving these issues will make an additional contribution to the prevention and treatment of osteoporosis, which is a global public health problem primarily in the elderly population. Also, it will contribute to the quality of life of people with bone injuries, osteoporosis or other bone diseases.

Acknowledgments

The authors would like to thank the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grants

No. 451-03-65/2024-03/200113 and 451-03-66/2024-03/20003) and the Faculty of Medicine, University of Niš for financial support provided by the Internal Project No. 68.

References

1. Mao W, Huang G, Chen H, Xu L, Qin S, Li A. Research progress of the role of anthocyanins on bone regeneration. *Front Pharmacol* 2021; 12:773660. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Salhotra A, Shah HN, Levi B, Longaker MT. Mechanisms of bone development and repair. *Nat Rev Mol Cell Biol* 2020;21(11):696-711. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Demontiero O, Vidal C, Duque G. Aging and bone loss: new insights for the clinician. *Ther Adv Musculoskelet Dis* 2012;4(2):61-76. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Simpson CR, Kelly HM, Murphy CM. Synergistic use of biomaterials and licensed therapeutics to manipulate bone remodelling and promote non-union fracture repair. *Adv Drug Deliv Rev* 2020; 160:212-33. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Mende W, Götzl R, Kubo Y, Pufe T, Ruhl T, Beier JP. The role of adipose stem cells in bone regeneration and bone tissue engineering. *Cells* 2021;10(5):975. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Baldwin P, Li DJ, Auston DA, Mir HS, Yoon RS, Koval KJ. Autograft, allograft, and bone graft substitutes: clinical evidence and indications for use in the setting of orthopaedic trauma surgery. *J Orthop Trauma* 2019;33(4):203-13. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Barba M, Di Taranto G, Lattanzi W. Adipose-derived stem cell therapies for bone regeneration. *Expert Opin Biol Ther* 2017;17(6):677-89. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Bishop AT, Pelzer M. Vascularized bone allotransplantation: current state and implications for future reconstructive surgery. *Orthop Clin North Am* 2007;38(1):109-22. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Ansari M. Bone tissue regeneration: biology, strategies and interface studies. *Prog Biomater* 2019;8(4):223-37. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Healy KE, Guldberg RE. Bone tissue engineering. *J Musculoskelet Neuronal Interact* 2007;7(4):328-30. [\[PubMed\]](#)
11. Cvetković VJ, Najdanović JG, Vukelić-Nikolić MĐ, Stojanović S, Najman SJ. Osteogenic potential of *in vitro* osteo-induced adipose-derived mesenchymal stem cells combined with platelet-rich plasma in an ectopic model. *Int Orthop* 2015;39(11):2173-80. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Najdanović JG, Cvetković VJ, Stojanović S, Vukelić-Nikolić MĐ, Stanisavljević MN, Živković JM, et al. The influence of adipose-derived stem cells induced into endothelial cells on ectopic vasculogenesis and osteogenesis. *Cell Mol Bioeng* 2015;8(4):577-90. [\[CrossRef\]](#)
13. Živković JM, Najman SJ, Vukelić MĐ, Stojanović S, Aleksić M, Stanisavljević M, et al. Osteogenic effect of inflammatory macrophages loaded onto mineral bone substitute in subcutaneous implants. *Arch Biol Sci* 2015;67(1):173-86. [\[CrossRef\]](#)
14. Vukelić-Nikolić MĐ, Najman SJ, Vasiljević PJ, Jevtović-Stoimenov TM, Cvetković VJ, Andrejev MN, et al. Osteogenic capacity of diluted platelet-rich plasma in ectopic bone-forming model: Benefits for bone regeneration. *J Craniomaxillofac Surg* 2018;46(11):1911-8. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Barbeck M, Najman S, Stojanović S, Mitić Ž, Živković JM, Choukroun J, et al. Addition of blood to a phycogenic bone substitute leads to increased *in vivo* vascularization. *Biomed Mater* 2015;10(5):055007. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Živković JM, Stojanović ST, Vukelić-Nikolić MĐ, Radenković MB, Najdanović JG, Ćirić M, et al. Macrophages' contribution to ectopic osteogenesis in combination with blood clot and bone substitute: possibility for application in bone regeneration strategies. *Int Orthop* 2021;45(4):1087-95. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Najman SJ, Cvetković VJ, Najdanović JG, Stojanović S, Vukelić-Nikolić MĐ, Vučković I, et al. Ectopic osteogenic capacity of freshly isolated adipose-derived stromal vascular fraction cells supported with platelet-rich plasma: A simulation of intraoperative procedure. *J Craniomaxillofac Surg* 2016;44(10):1750-60. [\[CrossRef\]](#) [\[PubMed\]](#)
18. Najdanović JG, Cvetković VJ, Stojanović ST, Vukelić-Nikolić MĐ, Živković JM, Najman SJ. Vascularization and osteogenesis in ectopically implanted bone tissue-engineered constructs with endothelial and osteogenic differentiated adipose-derived stem cells. *World J Stem Cells* 2021;13(1):91-114. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Shimizu S, Matsushita H, Morii Y, Ohyama Y, Morita N, Tachibana R, et al. Effect of anthocyanin-rich bilberry extract on bone metabolism in ovariectomized rats. *Biomed Rep* 2018;8(2):198-204. [\[CrossRef\]](#) [\[PubMed\]](#)
20. Almeida M, O'Brien CA. Basic biology of skeletal aging: role of stress response pathways. *J Gerontol A Biol Sci Med Sci* 2013;68(10):1197-208. [\[CrossRef\]](#) [\[PubMed\]](#)
21. Carochi M, Ferreira IC. A review on antioxidants, prooxidants and related controversy: natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food Chem Toxicol* 2013; 51:15-25. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Bai XC, Lu D, Bai J, Zheng H, Ke ZY, Li XM, et al. Oxidative stress inhibits osteoblastic

- differentiation of bone cells by ERK and NF-kappaB. *Biochem Biophys Res Commun* 2004;314(1):197-207. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Domazetovic V, Marcucci G, Iantomasi T, Brandi ML, Vincenzini MT. Oxidative stress in bone remodeling: role of antioxidants. *Clin Cases Miner Bone Metab* 2017;14(2):209-16. [\[CrossRef\]](#) [\[PubMed\]](#)
 24. Joshi DD, Deb L, Somkuwar BG, Rana VS. Potential use of barks of woody vascular plants in bone mending: A review. *Saudi Pharm J* 2023;31(9):101714. [\[CrossRef\]](#) [\[PubMed\]](#)
 25. Suneetha J, Prasanthi S, Ramarao N, Seetharami BVA, Reddi TVV. Indigenous phytotherapy for bone fractures from Eastern Ghats. *Indian J Tradit Knowl* 2011;10(3):550-3.
 26. Sharma C, Kumari T, Arya KR. Ethnopharmacological survey on bone healing plants with special references to *Pholidota articulata* and *Coelogyne cristata* (Orchidaceae) used in folk tradition of Kumaon Uttarkhand India. *Int J Pharma Res Health Sci* 2014;2:185-90.
 27. Hardcastle AC, Aucott L, Fraser WD, Reid DM, Macdonald HM. Dietary patterns, bone resorption and bone mineral density in early post-menopausal Scottish women. *Eur J Clin Nutr* 2011;65(3):378-85. [\[CrossRef\]](#) [\[PubMed\]](#)
 28. Đudarić L, Fužinac-Smojver A, Muhvić D, Giacometti J. The role of polyphenols on bone metabolism in osteoporosis. *Food Research International* 2015;77(2):290-8. [\[CrossRef\]](#)
 29. He J, Giusti MM. Anthocyanins: natural colorants with health-promoting properties. *Annu Rev Food Sci Technol* 2010;1:163-87. [\[CrossRef\]](#) [\[PubMed\]](#)
 30. Bilawal A, Ishfaq M, Gantumur M-A, Qayum A, Shi R, Fazilani SA, et al. A review of the bioactive ingredients of berries and their applications in curing diseases. *Food Biosci* 2021;44 (Part A): 101407. [\[CrossRef\]](#)
 31. Manach C, Scalbert A, Morand C, Rémésy C, Jiménez L. Polyphenols: food sources and bioavailability. *Am J Clin Nutr* 2004;79(5):727-47. [\[CrossRef\]](#) [\[PubMed\]](#)
 32. Wang Z, Wang D, Yang D, Zhen W, Zhang J, Peng S. The effect of icariin on bone metabolism and its potential clinical application. *Osteoporos Int* 2018;29(3):535-44. [\[CrossRef\]](#) [\[PubMed\]](#)
 33. Motohashi N, Sakagami H. Functionality of anthocyanins as alternative medicine. In: Motohashi N, editor. *Bioactive heterocycles VI. Topics in heterocyclic chemistry*. Berlin, Heidelberg: Springer; 2008. 15: p. 1-48. [\[CrossRef\]](#)
 34. Yang J, Zhang L, Ding Q, Zhang S, Sun S, Liu W, et al. Flavonoid-Loaded Biomaterials in Bone Defect Repair. *Molecules* 2023;28(19):6888. [\[CrossRef\]](#) [\[PubMed\]](#)
 35. Millar CL, Duclos Q, Blesso CN. Effects of dietary flavonoids on reverse cholesterol transport, HDL metabolism, and HDL function. *Adv Nutr* 2017;8(2):226-39. [\[CrossRef\]](#) [\[PubMed\]](#)
 36. Eker ME, Aaby K, Budic-Leto I, Brnčić SR, El SN, Karakaya S, et al. A review of factors affecting anthocyanin bioavailability: possible implications for the inter-individual variability. *Foods* 2019;9(1):2. [\[CrossRef\]](#) [\[PubMed\]](#)
 37. Wei Z, Yang H, Shi J, Duan Y, Wu W, Lyu L, et al. Effects of different light wavelengths on fruit quality and gene expression of anthocyanin biosynthesis in blueberry (*Vaccinium corymbosum*). *Cells* 2023;12(9):1225. [\[CrossRef\]](#) [\[PubMed\]](#)
 38. Liu Z, Liang T, Kang C. Molecular bases of strawberry fruit quality traits: Advances, challenges, and opportunities. *Plant Physiology* 2023;193(2):900-14. [\[CrossRef\]](#) [\[PubMed\]](#)
 39. Steyn WJ. Prevalence and functions of anthocyanins in fruits. In: Gould K, Davies K, Winefield C, editors. *Anthocyanins, biosynthesis, functions, and applications*. New York: Springer; 2009. p. 85-105 [\[CrossRef\]](#)
 40. An X, Tan T, Song Z, Guo X, Zhang X, Zhu Y, et al. Physiological response of anthocyanin synthesis to different light intensities in blueberry. *PLoS One* 2023;18(6):e0283284. [\[CrossRef\]](#) [\[PubMed\]](#)
 41. Rossi A, Serraino I, Dugo P, Paola RD, Mondello L, Genovese T, et al. Protective effects of anthocyanins from blackberry in a rat model of acute lung inflammation. *Free Radic Res* 2003;37(8):891-900. [\[CrossRef\]](#) [\[PubMed\]](#)
 42. Devareddy L, Hooshmand S, Collins JK, Lucas EA, Chai SC, Arjmandi BH. Blueberry prevents bone loss in ovariectomized rat model of postmenopausal osteoporosis. *J Nutr Biochem* 2008;19(10):694-9. [\[CrossRef\]](#) [\[PubMed\]](#)
 43. Leyva-López N, Gutierrez-Grijalva EP, Ambríz-Pérez DL, Heredia JB. Flavonoids as cytokine modulators: a possible therapy for inflammation-related diseases. *Int J Mol Sci* 2016;17(6):921. [\[CrossRef\]](#) [\[PubMed\]](#)
 44. Thilavech T, Ngamukote S, Belobrajdic D, Abeywardena M, Adisakwattana S. Cyanidin-3-rutinoside attenuates methylglyoxal-induced protein glycation and DNA damage via carbonyl trapping ability and scavenging reactive oxygen species. *BMC Complement Altern Med* 2016;16:138. [\[CrossRef\]](#) [\[PubMed\]](#)
 45. Tabart J, Kevers C, Evers D, Dommès J. Ascorbic acid, phenolic acid, flavonoid, and carotenoid profiles of selected extracts from *Ribes nigrum*. *J Agric Food Chem* 2011;59(9):4763-70. [\[CrossRef\]](#) [\[PubMed\]](#)
 46. Scalzo J, Currie A, Stephens J, McGhie T, Alspach P, Horticulture and food research institute of New Zealand limited hortresearch. The anthocyanin composition of different *Vaccinium*, *Ribes* and *Rubus* genotypes. *Biofactors* 2008;34(1):13-21. [\[CrossRef\]](#) [\[PubMed\]](#)
 47. Gopalan A, Reuben SC, Ahmed S, Darvesh AS, Hohmann J, Bishayee A. The health benefits of blackcurrants. *Food Funct* 2012;3(8): 795-809. [\[CrossRef\]](#) [\[PubMed\]](#)
 48. Lee SG, Kim B, Yang Y, Pham TX, Park YK, Manatou J, et al. Berry anthocyanins suppress the expression and secretion of proinflammatory mediators in macrophages by inhibiting nuclear translocation of NF-κB independent of NRF2-mediated mechanism. *J Nutr Biochem* 2014;25(4), 404-11. [\[CrossRef\]](#) [\[PubMed\]](#)
 49. Lee SG, Vance TM, Nam TG, Kim DO, Koo SI, Chun OK. Contribution of anthocyanin composition to total antioxidant capacity of berries. *Plant Foods Hum Nutr* 2015;70(4):427-32. [\[CrossRef\]](#) [\[PubMed\]](#)
 50. Noda Y, Kaneyuki T, Mori A, Packer L. Antioxidant activities of pomegranate fruit extract and its anthocyanidins: delphinidin, cyanidin, and pelargonidin. *J Agric Food Chem* 2002;50(1):166-71. [\[CrossRef\]](#) [\[PubMed\]](#)
 51. Yang W, Guo Y, Liu M, Chen X, Xiao X, Wang S, et al. Structure and function of blueberry anthocyanins: A review of recent advances. *J Funct Foods* 2022;88:104864. [\[CrossRef\]](#)

52. He F, Mu L, Yan GL, Liang NN, Pan QH, Wang J, et al. Biosynthesis of anthocyanins and their regulation in colored grapes. *Molecules* 2010;15(12):9057-91. [\[CrossRef\]](#) [\[PubMed\]](#)
53. Vega EN, García-Herrera P, Ciudad-Mulero M, Dias MI, Matallana-González MC, Cámara M, et al. Wild sweet cherry, strawberry and bilberry as underestimated sources of natural colorants and bioactive compounds with functional properties. *Food Chemistry* 2023;414:135669. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Xue L, Otieno M, Colson K, Neto C. Influence of the growing region on the phytochemical composition and antioxidant properties of north american cranberry fruit (*Vaccinium macrocarpon* Aiton). *Plants* 2023;12(20):3595. [\[CrossRef\]](#) [\[PubMed\]](#)
55. Teng H, Fang T, Lin Q, Song H, Liu B, Chen L. Red raspberry and its anthocyanins: Bioactivity beyond antioxidant capacity. *Trends Food Sci Technol* 2017;66:153-65. [\[CrossRef\]](#)
56. Djordjević B, Šavikin K, Zdunić G, Janković T, Vulić T, Oparnica Č, et al. Biochemical properties of red currant varieties in relation to storage. *Plant Foods Hum Nutr* 2010;65:326-32. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Speer H, D'Cunha NM, Alexopoulos NI, McKune AJ, Naumovski N. Anthocyanins and human health—a focus on oxidative stress, inflammation and disease. *Antioxidants (Basel)* 2020;9(5):366. [\[CrossRef\]](#) [\[PubMed\]](#)
58. Moriwaki S, Suzuki K, Muramatsu M, Nomura A, Inoue F, Into T, et al. Delphinidin, one of the major anthocyanidins, prevents bone loss through the inhibition of excessive osteoclastogenesis in osteoporosis model mice. *PLoS One* 2014;9(5):e97177. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Maggio D, Barabani M, Pierandrei M, Polidori MC, Catani M, Mecocci P, et al. Marked decrease in plasma antioxidants in aged osteoporotic women: results of a cross-sectional study. *J Clin Endocrinol Metab* 2003;88(4):1523-7. [\[CrossRef\]](#) [\[PubMed\]](#)
60. Banfi G, Iorio EL, Corsi MM. Oxidative stress, free radicals and bone remodeling. *Clin Chem Lab Med* 2008;46(11):1550-5. [\[CrossRef\]](#) [\[PubMed\]](#)
61. Dreher I, Schütze N, Baur A, Hesse K, Schneider D, Köhrle J, et al. Selenoproteins are expressed in fetal human osteoblast-like cells. *Biochem Biophys Res Commun* 1998;245(1):101-7. [\[CrossRef\]](#) [\[PubMed\]](#)
62. Zhai ZJ, Li HW, Liu GW, Qu XH, Tian B, Yan W, et al. Andrographolide suppresses RANKL-induced osteoclastogenesis *in vitro* and prevents inflammatory bone loss *in vivo*. *Br J Pharmacol* 2014;171(3):663-75. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Domazetovic V, Marcucci G, Falsetti I, Bilia AR, Vincenzini MT, Brandi ML, et al. Blueberry juice antioxidants protect osteogenic activity against oxidative stress and improve long-term activation of the mineralization process in human osteoblast-like SaOS-2 Cells: Involvement of SIRT1. *Antioxidants (Basel)* 2020;9(2):125. [\[CrossRef\]](#) [\[PubMed\]](#)
64. Liu J, Han W, Chen L, Tang K. Mechanism of osteogenic and adipogenic differentiation of tendon stem cells induced by sirtuin 1. *Mol Med Rep* 2016;14(2):1643-8. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Ostos Mendoza KC, Garay Buenrostro KD, Kanabar PN, Maienschein-Cline M, Los NS, Arbieva Z, et al. Peonidin-3-O-glucoside and resveratrol increase the viability of cultured human hFOB Osteoblasts and alter the expression of genes associated with apoptosis, osteoblast differentiation and osteoclastogenesis. *Nutrients* 2023;15(14):3233. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Nagaoka M, Maeda T, Chatani M, Handa K, Yamakawa T, Kiyohara S, et al. A delphinidin-enriched maqui berry extract improves bone metabolism and protects against bone loss in osteopenic mouse models. *Antioxidants (Basel)* 2019;8(9):386. [\[CrossRef\]](#) [\[PubMed\]](#)
67. Nagaoka M, Maeda T, Moriwaki S, Nomura A, Kato Y, Niida S, et al. Petunidin, a B-ring 5'-O-methylated derivative of delphinidin, stimulates osteoblastogenesis and reduces sRANKL-induced bone loss. *Int J Mol Sci* 2019;20(11):2795. [\[CrossRef\]](#) [\[PubMed\]](#)
68. Hu B, Chen L, Chen Y, Zhang Z, Wang X, Zhou B. Cyanidin-3-glucoside regulates osteoblast differentiation via the ERK1/2 signaling pathway. *ACS Omega* 2021;6(7):4759-66. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Kim JM, Yang YS, Park KH, Oh H, Greenblatt MB, Shim JH. The ERK MAPK pathway is essential for skeletal development and homeostasis. *Int J Mol Sci* 2019;20(8):1803. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Casati L, Pagani F, Fibiani M, Lo Scalzo R, Sibilia V. Potential of delphinidin-3-rutinoside extracted from *Solanum melongena* L. as promoter of osteoblastic MC3T3-E1 function and antagonist of oxidative damage. *Eur J Nutr* 2019;58(3):1019-32. [\[CrossRef\]](#) [\[PubMed\]](#)
71. Saulite L, Jekabsons K, Klavins M, Muceniece R, Riekstina U. Effects of malvidin, cyanidin and delphinidin on human adipose mesenchymal stem cell differentiation into adipocytes, chondrocytes and osteocytes. *Phytomedicine* 2019;53:86-95. [\[CrossRef\]](#) [\[PubMed\]](#)
72. Arjmandi BH, Lucas EA, Juma S, Soliman A, Stoecker BJ, Khalil DA, et al. Dried plums prevent ovariectomy-induced bone loss in rats. *JANA* 2001;4:50-6.
73. Li T, Wu SM, Xu ZY, Ou-Yang S. Rabbiteye blueberry prevents osteoporosis in ovariectomized rats. *J Orthop Surg Res* 2014;9:56. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Zheng X, Mun S, Lee SG, Vance TM, Hubert P, Koo SI, et al. Anthocyanin-rich blackcurrant extract attenuates ovariectomy-induced bone loss in mice. *J Med Food* 2016;19(4):390-7. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Sakaki J, Melough M, Lee SG, Kalinowski J, Koo SI, Lee SK, et al. Blackcurrant supplementation improves trabecular bone mass in young but not aged mice. *Nutrients* 2018;10(11):1671. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Lee SG, Kim B, Soung do Y, Vance T, Lee JS, Lee JY, et al. Relationship between oxidative stress and bone mass in obesity and effects of berry supplementation on bone remodeling in obese male mice: an exploratory study. *J Med Food* 2015;18(4):476-82. [\[CrossRef\]](#) [\[PubMed\]](#)
77. Zeng Z, Li H, Luo C, Hu W, Weng TJ, Shuang F. Pelargonidin ameliorates inflammatory response and cartilage degeneration in osteoarthritis via suppressing the NF-κB pathway. *Arch Biochem Biophys* 2023; 743:109668. [\[CrossRef\]](#) [\[PubMed\]](#)
78. Cheng J, Zhou L, Liu Q, Tickner J, Tan Z, Li X, et al. Cyanidin chloride inhibits ovariectomy-induced osteoporosis by suppressing RANKL-mediated osteoclastogenesis and associated signaling pathways. *J Cell Physiol* 2018;233(3):2502-12. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Dou C, Li J, Kang F, Cao Z, Yang X, Jiang H, et al. Dual effect of cyanidin on RANKL-induced

- differentiation and fusion of osteoclasts. *J Cell Physiol* 2016;231(3):558-67. [\[CrossRef\]](#) [\[PubMed\]](#)
80. Zhang J, Liu Z, Luo Y, Li X, Huang G, Chen H, et al. The role of flavonoids in the osteogenic differentiation of mesenchymal stem cells. *Front Pharmacol* 2022; 13:849513. [\[CrossRef\]](#) [\[PubMed\]](#)
 81. Welch A, MacGregor A, Jennings A, Fairweather-Tait S, Spector T, Cassidy A. Habitual flavonoid intakes are positively associated with bone mineral density in women. *J Bone Miner Res* 2012;27(9):1872-8. [\[CrossRef\]](#) [\[PubMed\]](#)
 82. Li JJ, Huang ZW, Wang RQ, Ma XM, Zhang ZQ, Liu Z, et al. Fruit and vegetable intake and bone mass in Chinese adolescents, young and postmenopausal women. *Public Health Nutr* 2012;16(1):78-86. [\[CrossRef\]](#) [\[PubMed\]](#)
 83. Zalloua PA, Hsu YH, Terwedow H, Zang T, Wu D, Tang G, et al. Impact of seafood and fruit consumption on bone mineral density. *Maturitas* 2007;56(1):1-11. [\[CrossRef\]](#) [\[PubMed\]](#)
 84. New SA, Robins SP, Campbell MK, Martin JC, Garton MJ, Bolton-Smith C, et al. Dietary influences on bone mass and bone metabolism: further evidence of a positive link between fruit and vegetable consumption and bone health? *Am J Clin Nutr* 2000;71(1):142-51. [\[CrossRef\]](#) [\[PubMed\]](#)

Pregledni rad

UDC: 612.753:547.973
doi: 10.5633/amm.2024.0412**BOBIČASTO VOĆE KAO PRIRODNO SREDSTVO
POTPORE KOSTIJU: UČEŠĆE ANTOCIJANA U
MOLEKULARNIM MEHANIZMIMA PROCESA ZALEČENJA
I REGENERACIJE***Jelena Najdanović^{1,2}, Stevo Najman^{1,2}, Jelena Živković^{1,2},
Bojana Miladinović³*¹Univerzitet u Nišu, Medicinski fakultet u Nišu, UNO Biologija sa humanom genetikom, Niš, Srbija²Univerzitet u Nišu, Medicinski fakultet u Nišu, Naučnoistraživački centar za biomedicinu,
Odeljenje za ćelijsko i tkivno inženjerstvo, Niš, Srbija³Univerzitet u Nišu, Medicinski fakultet u Nišu, Katedra za farmaciju, Niš, SrbijaKontakt: Jelena Najdanović
Bulevar dr Zorana Đinđića 81, 18000 Niš, Srbija
E-mail: jelena.najdanovic@medfak.ni.ac.rs

Koštano tkivo ima izuzetan potencijal samoregeneracije koji, u slučaju da dođe do povrede, omogućava povratak u potpuno funkcionalno stanje pre povrede. Međutim, starenje, bolesti, gojaznost, traume, višestruki prelomi, infekcije i uklanjanje tumora uzrokuju velike koštane defekte koji se ne mogu spontano zalečiti. Kako bi zaceljenje i regeneracija kostiju bili uspešni, razvijeni su različiti pristupi, među kojima su i primena autografta, alografta i tkivnog inženjerstva kosti. Jedan od pristupa zasniiva se na saznanju da gubitak kostiju kod ljudi i kod mnogih životinja u toku starenja delimično izaziva akumulacija reaktivnih vrsta kiseonika. Zbog spektra bioloških aktivnosti, uključujući i onu antioksidativnu, esencijalna polifenolna jedinjenja – antocijani – deo su značajne oblasti istraživanja povezane sa sredstvima i metodama koje se koriste za zaceljenje i regeneraciju kostiju. Bobičasto voće je posebno bogato antocijanima. Na osnovu *in vitro* i *in vivo* proučavanja molekularnih mehanizama uključenih u zaceljenje i regeneraciju kostiju koji su potpomognuti antocijanima iz bobičastog voća i na osnovu opservacionih istraživanja sprovedenih među ljudima, utvrđeno je da antocijani bobičastog voća pospešuju osteoblastogenezu, suzbijaju osteoklastogenezu i imaju osteoimunološku aktivnost. Dakle, antocijane iz bobičastog voća treba smatrati potencijalnim, široko rasprostranjenim terapijskim sredstvom davanja potpore kostima. Ipak, pre primene bobičastog voća kao prirodnog sredstva potpore kostima, treba se pozabaviti pojašnjenjem molekularnih mehanizama delovanja antocijana u metabolizmu kostiju i utvrđivanjem efikasnih doza konkretnih antocijana za terapiju regeneracije kostiju. Takođe, treba sprovesti kliničke studije kako bi se utvrdila terapijska efikasnost različitih tipova i koncentracija antocijana.

*Acta Medica Medianae 2024; 63(4): 104–113.***Ključne reči:** zaceljenje kostiju, regeneracija kostiju, bobičasto voće, antocijani, molekularni mehanizmi

"This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) Licence".