

Original article
DOI: 10.5633/amm.2024.0203

COVID-19 AND IMMUNOTHERAPY IN JUVENILE IDIOPATHIC ARTHRITIS

Nikola Jovanović¹, Milica Jordanović², Jelena Vojinović³, Milena Manojlović³

¹ General Hospital Bor, Pediatrics department, Bor, Serbia

² Healthcare center Užice, Užice, Serbia

³ University Clinical Centar Niš, Pediatric Clinic, Niš, Serbia

Contact: Nikola Jovanović
Dragiše Cvetkovića 23 str., 18000 Niš, Serbia
Email: nikola92.jovanovic92@gmail.com
Tel: 0638883247

AMM Paper Accepted

Abstract

The COVID-19 pandemic was a huge challenge to pediatricians around the world. Patients with juvenile idiopathic arthritis (JIA) and COVID-19 infection could present a potential ethical and medical dilemma. Here we discuss the results of a medical survey made on the parents of children with JIA. Our primary aim was to determine if there was a significant difference in the number of flares of JIA after COVID-19 infection between the group of children who were receiving biological drugs and children not receiving biologicals. Other goals were to investigate the parents' motivation to vaccinate children against SARS-CoV-2 and to determine the most frequent symptoms of COVID-19 infection in these children. A retrospective study was based on the data of a telephone survey conducted between March 10, 2022 and May 12, 2022, including 65 pediatric patients with JIA. The data were provided from the "Heliant" information system database and survey results. In children who tested positive for SARS-CoV-2, the most frequent symptom was fever, followed by the upper respiratory symptoms. Four flares of JIA were observed in the group of children on biological therapy, while in the group without biologicals there were two flares that followed the COVID-19 infection. The parents' motivation for vaccination against SARS-CoV-2 was extremely low. Our survey-based research did not find a significant difference regarding the COVID-19 infection between children with JIA on biologicals and children with JIA not receiving biologicals, but it did emphasize the parents' hesitancy about vaccination. We propose building a unique database for patients with the diagnosis of JIA which could improve the quality of life of these patients.

Keywords: COVID-19, JIA, SARS-CoV-2, flare, biologicals

Originalni rad
DOI: 10.5633/amm.2024.0203

KOVID-19 I IMUNOTERAPIJA U JUVENILNOM IDIOPATSKOM ARTRITISU

Nikola Jovanović¹, Milica Jordanović², Jelena Vojinović³, Milena Manojlović³

¹Opšta bolnica Bor, Odeljenje pedijatrije, Bor, Srbija

²Dom zdravlja Užice, Užice, Srbija

³Univerzitetski klinički centar Niš, Klinika za pedijatriju, Niš, Srbija

Kontakt: Nikola Jovanović
Dragiše Cvetkovića 23, Niš
Email: nikola92.jovanovic92@gmail.com
Tel: 0638883247

Apstrakt

Pandemija Kovida-19 predstavlja ogroman izazov pedijatrijama širom sveta. Deca obolela od juvenilnog idiopatskog artritisa (JIA) koja imaju Kovid-19 i koja su po indikacijama na biološkoj terapiji, mogu predstavljati medicinsku i etičku dilemu. U ovom radu nastojali smo da odgovorimo na pitanje da li postoji povećana sklonost ka egzacerbacijama JIA nakon preležane infekcije Kovid-19 kod dece sa JIA na biološkoj terapiji u odnosu na decu sa JIA koja nisu na biološkoj terapiji. Pored toga, ispitali smo i motivaciju roditelja za vakcinaciju dece i simptome bolesti Kovid-19 u ovoj populaciji. Urađena je retrospektivna studija, na osnovu rezultata ankete rađene između 10.03.2022 i 12.05.2022 koja je obuhvatila 65-oro pedijatrijskih pacijenata. Anketa je sadržala niz zatvorenih i otvorenih pitanja, na koja su roditelji davali odgovore. Jedan deo podataka, koji se ticao trajanja JIA, brojeva telefona i terapije preuzet je iz "Heliant" informacionog sistema. Grupe su poredene putem Hi-kvadrat testa. Najčešći simptom bila je temperatura, a potom simptomi gornjeg respiratornog trakta. U četiri slučaja zapažena je egzacerbacija osnovne bolesti u grupi dece na biološkoj terapiji sa JIA, koje je usledilo nakon Kovida-19, dok je u grupi dece sa JIA bez biološke terapije zabeleženo dva pogoršanja. Između ove dve grupe nije nađeno značajno statističko odstupanje. Motivacija roditelja za vakcinaciju protiv SARS-CoV-2 bila je izuzetno niska. Studija na većem broju pacijenta sa JIA može dati pouzdanije podatke o uticaju Kovida-19 na ovu grupu pacijenata. Predlažemo formiranje jedinstvene baze podataka za ove pacijente, koja kliničarima u budućnosti može pomoći u poboljšanju kvaliteta života dece sa JIA.

Ključne reči: JIA, Kovid-19, SARS-CoV-2, biološka terapija, egzacerbacija.

Introduction

The start of COVID-19 pandemic was pronounced on March 11, 2020 by the World Health Organization (WHO). From that moment, until writing the introductory part of this article, it was estimated that the disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) took approximately 20 million lives (1), thus leaving deep and unfathomable consequences on society, medicine, economy, however, it also contributed to the development of various sociological and cultural phenomena. SARS-CoV-2 is an infective agent constituted of nucleocapsid, membrane, envelope, and spike (S) protein. It is characterized by a high transmission efficacy and genetic similarity to SARS and MERS viruses. The spike protein engages with ACE-2 receptors of human cells, which allows an entry of a virus particle in the cell (2-5). Juvenile idiopathic arthritis is the most common chronic rheumatic disease in children. Its etiology is still unknown, although it is thought to be provoked by a complex interaction between genetic and environmental factors such as infections, human leukocyte antigen (HLA), gut microbiota, etc. It is classified by The International League of Associations for Rheumatology (ILAR) into several subtypes (6-9). Due to its pathophysiology, it is considered a polygenic autoinflammatory disease. Disease-modifying antirheumatic drugs (DMARDs) represent a group of drugs used in the treatment of rheumatic diseases in pediatric and adult population. A specific mechanism of monoclonal antibodies provides targeted neutralization of certain cytokines involved in the pathogenesis of JIA. These medications can silence the disease course and provide remission periods for patients with JIA (10, 11). Although interleukin-6, interleukin-1 and tumor necrosis factor inhibitors safety profile are regarded highly acceptable, their use in patients with JIA, and especially in patients with comorbidities, could provoke an increased risk of infections (12-14). Another aspect of COVID-19 pandemic is dealing with covid pediatric patients that have comorbidities. One such population are patients with JIA. Pfizer-BioNTech COVID-19 vaccine is an mRNA-based vaccine and the only vaccine registered for use in pediatric population in Serbia. Vaccine hesitancy, mostly caused by misinformation, was a major problem in fighting with COVID-19 pandemic, especially in pediatric patients (15). Potential flares that can be provoked by other infections could possibly occur after COVID-19 infection. Fever that can occur in both covid and JIA must be carefully evaluated. Other important aspects are when to vaccinate, taking in consideration immunosuppressive therapy in JIA patients. Therefore, patients with JIA and COVID-19 infection could present a potential ethical and medical dilemma, and every single case must be closely monitored. Here we discuss the results of a medical survey conducted during pandemics, involving parents whose children were diagnosed with JIA.

Aim

Our primary aim was to determine if there was a significant difference in the number of juvenile idiopathic arthritis flares that could be related to SARS-CoV-2 infection between the group of children with JIA who were receiving biological drugs and children with JIA not receiving biological drugs. Other goals were to investigate the

parents' motivation to vaccinate children against SARS-CoV-2 and to determine the most frequent symptoms of COVID-19 infection in this population.

Materials and methods

A retrospective study based on the data of a telephone survey was conducted between March 10, 2022 and May 12, 2022, and it included 65 pediatric patients diagnosed with juvenile idiopathic arthritis. Parents were beforehand informed to anticipate a call from a physician. The survey referred to the period from the beginning of the pandemic. Children were divided in two groups: one that was receiving monoclonal antibodies as a part of the treatment for JIA (n=39), and the control group that consisted of children with the diagnosis of JIA, who were not receiving biological drugs. The control group (n=26) included children who were receiving methotrexate and other DMARDs, but also children in disease remission. Some of the research data considering the total duration of JIA, parent's phone numbers and treatment was provided through the "Heliant" database. Other data was provided from the survey. The survey contained a series of closed and open-ended questions about COVID-19, to which parents gave answers, and it could be roughly divided into four clusters: 1) Positivity or prolonged contact with household members; 2) Symptoms that predominated infection; 3) Potential flares of JIA that followed the infection; 4) Vaccinal status or parents motivation to vaccinate children against SARS-CoV-2. A Chi-squared test was used for group comparison.

Results

Gender, age, and duration of the disease (JIA) are shown in Table 1. The number of positive tested children or children with JIA in prolonged contact with household members tested positive for COVID-19, number of flares that followed COVID-19 infection and number of parents that vaccinated or were motivated for vaccination of their children are shown in Table 2, along with the most frequently noted symptoms. Therapy of children on biological drugs was adalimumab (n=18, 48,6%), etanercept (n=11, 29,7%) and tocilizumab (n=8, 21,6%) together with methotrexate. For two cases, the data for the type of biological treatment in "Heliant" healthcare information system database were not available. In the group without biologicals, treatment was methotrexate (n=14, 53,8%), hydroxychloroquine (n=2, 7,7%), naproxen with methotrexate (n=2, 7,7%), naproxen (n=2, 7,7%), prednisone with methotrexate (n=1, 3,8%) and physical therapy (n=5, 19,2%). In one case (n=1, 3,84%), the disease exacerbation occurred after parents did not show up for regular check-ups during lockdown for about two months. In children that tested positive for SARS-CoV-2 on PCR or antigen test (n=22), the most frequent symptom was fever (n=14, 63,6%). Fever was <39 °C in all cases, except in one where it measured above 39 °C (on one occasion 39,8 °C). Other symptoms noted were the upper respiratory symptoms (cough, nasal congestion, sore throat, etc.), fatigue, myalgias and conjunctivitis. Parents of three children noted an asymptomatic infection (n=3, 14%) that was proven with antibody testing. In most cases, symptoms were mild and mostly lasted less than a week. In one case, parents noted that their child had loss of smell and taste problems that persisted for almost six months. No significant statistical difference was found in the number of flares that followed COVID-19 infection between children with JIA who were receiving biological drugs and children with the diagnosis of JIA not receiving biological drugs (χ^2 with Yates correction was 0.0077, p= .930299. Not significant at p < .05). As a reason for not vaccinating their children or lack of motivation for vaccination, most parents reported a misconception that vaccines were "not tested enough and are experimental". Relative risk

for flares was $R=1.33$ ($R>1$) in the group of children receiving biologicals compared to the group of children without biologicals in their treatment.

Table 1. Characteristics of JIA patients

Gender	Biological therapy		Control group		Total
	Male	Female	Male	Female	Male and female
	10 (25,64%)	29 (74,35%)	11 (40,74%)	15 (59,25%)	65 (100%)
Age	11.75±4.70		9.71±4.61		10.87±4.75
Disease duration*	7.11±10.90 [†]		3.93±2.32 [†]		5.84±8.67

*Total time with the diagnosis of JIA. [†]Disease duration was not known for one case for biological therapy and two cases in the control group. Biological therapy group showed greater variation of data for disease duration.

Table 2. Survey results

	Biological therapy	Control group	Total	Symptoms
Positive/contact*	31 (79,48%)	20 (76,93%)	51 (78,46%)	-Fever -Nasal congestion, rhinorrhoea
Flares of JIA[†]	4 (10,25%)	2 (7,70%)	7 (10,78%) [‡]	-Cough, sore throat
Motivated/vaccinated	3 (7,69%)	1 (3,84%)	4 (6,15%)	-Myalgias, fatigue

*Tested positive or in prolonged contact with family or household members who tested positive on COVID-19. [†]Flares of JIA that followed COVID-19 infection. [‡]In one case, exacerbation of JIA did not occur after covid infection. No significant statistical difference b/w the groups was noted $p>0.05$.

Discussion

In our survey-based research, we tried to examine whether there was a significant difference in the number of flares that could be related to SARS-CoV-2 infection, between pediatric patients with the diagnosis of JIA who were on biological therapy and pediatric patients with the diagnosis of JIA not receiving these drugs.

Key points:

- Patients with the diagnosis of juvenile idiopathic arthritis (JIA) usually had a mild form of COVID-19 disease.
- There were cases where flares of JIA occurred after the infection with SARS-CoV-2.
- We failed to prove that greater risk of flares after SARS-CoV-2 infection existed in the group of pediatric patients with JIA who were receiving biological drugs.
- It was evident that parents' motivation to vaccinate their children against SARS-CoV-29 was extremely low.

The course of COVID-19 infection in children is usually milder and more asymptomatic compared to adult population. Most frequent symptoms noted by authors are fever, cough, and rhinorrhoea, which is consistent with our results (Table 2). Most pediatric patients who required ICU admission had comorbidities such as cancer (16-17). The course of COVID-19 infection in pediatric patients with JIA is generally not much different than the course of infection in the rest of pediatric population, although the risk of exacerbations of JIA triggered by infection was observed (18). This was also evident in our study. It was noted that COVID-19 infection could provoke flares of JIA in pediatric patients in remission or with inactive disease (19, 20). Increased relapse rate during lockdown was found in one Italian study (21), which could be caused by organization problems and general healthcare management. Nevertheless, the full impact of COVID-19 is still to be investigated for this group of patients. The total number of disease exacerbations observed in our study could also be caused by factors other than COVID-19, not evaluated by the survey (prolonged period off drugs, worsening of disease activity, stress, etc.). We could speculate about a protective effect of TNF- α blockers against cytokine storm syndrome (CSS) in pediatric patients with JIA who are on TNF- α inhibitors, as this potential effect was described in adult patients with rheumatic disease (22, 23). A small discrepancy between our two groups in the number of flares after COVID-19 (biological vs. without biologicals, 4:2, with RR=1.33) could be a consequence of disease severity, however, a larger study is needed for evaluating the true meaning of these data. Patients that required monoclonal antibodies in treatment are more likely to have more serious illness and likely more prone to disease exacerbations. Smell and taste dysfunction that lasted for almost six months was noted in one child in our study. Prolonged loss of taste and smell is a symptom described as part of "long COVID" (24, 25). Parents of children who were receiving biological drugs talked more freely about COVID-19 vaccination, and there is a slight discrepancy in motivation towards vaccination between the two groups (biological vs. without biologicals, 3:1). This could be caused by frequent contact with healthcare workers. Parents' extremely low motivation for vaccinating their children against COVID-19 can be caused by multiple factors. A significant component of vaccination hesitancy may be a conflicting and inadequate media coverage (26, 27). Parents who are vaccinated are more likely to vaccinate their children. The important factor for this decision may be parents' level of education (28, 29). Having in mind that most parents noted as a reason for not vaccinating their children that vaccines were "not tested enough and are experimental", an adequate strategy and promotion campaign could present a potential solution for this problem (30, 31).

Conclusion

Our survey-based research did not demonstrate a significant difference in terms of COVID-19 infection-provoked flares between children with JIA who were receiving biological drugs and children with JIA who were not on biologicals, but it did emphasize

the problems with parents' trust in vaccines, so similar surveys can be helpful in the future, and on the larger scale they could provide more significant and reliable data. Therefore, we propose building a unique database for pediatric patients with the diagnosis of juvenile idiopathic arthritis, so that healthcare workers dealing with this population could have a better concept on how to provide better quality of life for their patients.

Reference

1. Wise J. Covid-19: Global death toll may be three times higher than official records, study suggests. *BMJ*. Published online March 10, 2022:o636. doi:10.1136/bmj.o636

2. Kirtipal N, Bharadwaj S, Kang SG. From SARS to SARS-CoV-2, insights on structure, pathogenicity and immunity aspects of pandemic human coronaviruses. *Infect Genet Evol* 2020;85:104502. doi:10.1016/j.meegid.2020.104502
3. Jackson CB, Farzan M, Chen B, Choe H. Mechanisms of SARS-CoV-2 entry into cells. *Nat Rev Mol Cell Biol* 2022;23(1):3-20. doi:10.1038/s41580-021-00418-x
4. Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19 [published correction appears in *Nat Rev Microbiol*. 2022 Feb 23;:]. *Nat Rev Microbiol* 2021;19(3):141-154. doi:10.1038/s41579-020-00459-7
5. Xu X, Chen P, Wang J, et al. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci China Life Sci* 2020;63(3):457-460. doi:10.1007/s11427-020-1637-5.
6. Barut K, Adrovic A, Şahin S, Kasapçopur Ö. Juvenile Idiopathic Arthritis. *Balkan Med J* 2017;34(2):90-101. doi:10.4274/balkanmedj.2017.0111
7. Verwoerd A, Ter Haar NM, de Roock S, Vastert SJ, Bogaert D. The human microbiome and juvenile idiopathic arthritis. *Pediatr Rheumatol Online J* 2016;14(1):55. Published 2016 Sep 20. doi:10.1186/s12969-016-0114-4
8. Lee JJY, Schneider R. Systemic Juvenile Idiopathic Arthritis. *Pediatr Clin North Am* 2018;65(4):691-709. doi:10.1016/j.pcl.2018.04.005
9. Prakken B, Albani S, Martini A. Juvenile idiopathic arthritis. *Lancet* 2011;377(9783):2138-2149. doi:10.1016/S0140-6736(11)60244-4
10. Welzel T, Winskill C, Zhang N, Woerner A, Pfister M. Biologic disease modifying antirheumatic drugs and Janus kinase inhibitors in paediatric rheumatology - what we know and what we do not know from randomized controlled trials [published

- correction appears in *Pediatr Rheumatol Online J*. 2021 Aug 16;19(1):124]. *Pediatr Rheumatol Online J* 2021;19(1):46. Published 2021 Mar 25. doi:10.1186/s12969-021-00514-4
11. Cimaz R, Maioli G, Calabrese G. Current and emerging biologics for the treatment of juvenile idiopathic arthritis. *Expert Opin Biol Ther* 2020;20(7):725-740. doi:10.1080/14712598.2020.1733524
 12. Thiele F, Klein A, Windschall D, et al. Comparative risk of infections among real-world users of biologics for juvenile idiopathic arthritis: data from the German BIKER registry. *Rheumatol Int* 2021;41(4):751-762. doi:10.1007/s00296-020-04774-3
 13. Winthrop KL, Chiller T. Preventing and treating biologic-associated opportunistic infections. *Nat Rev Rheumatol* 2009;5(7):405-410. doi:10.1038/nrrheum.2009.105
 14. Giles JT, Bathon JM. Serious infections associated with anticytokine therapies in the rheumatic diseases. *J Intensive Care Med* 2004;19(6):320-334. doi:10.1177/0885066604267854
 15. Evans S, Klas A, Mikocka-Walus A, et al. "Poison" or "protection"? A mixed methods exploration of Australian parents' COVID-19 vaccination intentions. *J Psychosom Res* 2021;150:110626. doi:10.1016/j.jpsychores.2021.110626
 16. Qi K, Zeng W, Ye M, et al. Clinical, laboratory, and imaging features of pediatric COVID-19: A systematic review and meta-analysis. *Medicine (Baltimore)* 2021;100(15):e25230. doi:10.1097/MD.00000000000025230
 17. Alsohime F, Temsah MH, Al-Nemri AM, Somily AM, Al-Subaie S. COVID-19 infection prevalence in pediatric population: Etiology, clinical presentation, and outcome. *J Infect Public Health* 2020;13(12):1791-1796. doi:10.1016/j.jiph.2020.10.008

18. Naddei R, Alfani R, Bove M, et al. Increased relapse rate during COVID-19 lockdown in an Italian cohort of children with juvenile idiopathic arthritis [published online ahead of print, 2021 Aug 16]. *Arthritis Care Res (Hoboken)* 2021;10.1002/acr.24768. doi:10.1002/acr.24768
19. Hügler B, Krumrey-Langkammerer M, Haas JP. Infection with SARS-CoV-2 causes flares in patients with juvenile idiopathic arthritis in remission or inactive disease on medication. *Pediatr Rheumatol Online J* 2021;19(1):163. Published 2021 Nov 29. doi:10.1186/s12969-021-00653-8
20. Fernández-Sarmiento J, De Souza D, Jabornisky R, Gonzalez GA, Arias López MDP, Palacio G. Paediatric inflammatory multisystem syndrome temporally associated with COVID-19 (PIMS-TS): a narrative review and the viewpoint of the Latin American Society of Pediatric Intensive Care (SLACIP) Sepsis Committee. *BMJ Paediatr Open* 2021;5(1):e000894. Published 2021 Feb 4. doi:10.1136/bmjpo-2020-000894
21. Naddei R, Alfani R, Bove M, et al. Increased relapse rate during COVID-19 lockdown in an Italian cohort of children with juvenile idiopathic arthritis [published online ahead of print, 2021 Aug 16]. *Arthritis Care Res (Hoboken)* 2021;10.1002/acr.24768. doi:10.1002/acr.24768
22. Tripathi K, Godoy Brewer G, Thu Nguyen M, et al. COVID-19 and Outcomes in Patients With Inflammatory Bowel Disease: Systematic Review and Meta-Analysis [published online ahead of print, 2021 Oct 27]. *Inflamm Bowel Dis* 2021. doi:10.1093/ibd/izab236
23. Salesi M, Shojaie B, Farajzadegan Z, Salesi N, Mohammadi E. TNF- α Blockers Showed Prophylactic Effects in Preventing COVID-19 in Patients with Rheumatoid Arthritis and Seronegative Spondyloarthropathies: A Case-Control Study

- [published online ahead of print, 2021 Jul 23]. *Rheumatol Ther* 2021;1-16.
doi:10.1007/s40744-021-00342-8
24. Borch L, Holm M, Knudsen M, Ellermann-Eriksen S, Hagstroem S. Long COVID symptoms and duration in SARS-CoV-2 positive children - a nationwide cohort study. *Eur J Pediatr* 2022;181(4):1597-1607. doi:10.1007/s00431-021-04345-z
25. Elmas B, Çavdaroglu PD, Orhan MF, et al. Evaluation of taste and smell disorders in pediatric COVID-19 Cases. *Rev Assoc Med Bras (1992)* 2021;67(6):789-794. doi:10.1590/1806-9282.20200547
26. Thunström L, Ashworth M, Finnoff D, Newbold SC. Hesitancy Toward a COVID-19 Vaccine. *Ecohealth* 2021;18(1):44-60. doi:10.1007/s10393-021-01524-0
27. Paris C, Bénézit F, Geslin M, et al. COVID-19 vaccine hesitancy among healthcare workers. *Infect Dis Now* 2021;51(5):484-487. doi:10.1016/j.idnow.2021.04.001
28. Naso J, Rojas S, Peng J, et al. High Parental Vaccine Motivation at a Neighborhood-Based Vaccine and Testing Site Serving a Predominantly Latinx Community. *Health Equity* 2021;5(1):840-846. doi:10.1089/heq.2021.0073
29. Pan F, Zhao H, Nicholas S, Maitland E, Liu R, Hou Q. Parents' Decisions to Vaccinate Children against COVID-19: A Scoping Review. *Vaccines (Basel)* 2021;9(12):1476. doi:10.3390/vaccines9121476
30. Barello S, Maiorino G, Palamenghi L, et al. Exploring the Motivational Roots of Getting Vaccinated against COVID-19 in a Population of Vaccinated Pediatric Healthcare Professionals: Evidence from an Italian Cross-Sectional Study. *Vaccines (Basel)* 2022;10(3):467. doi:10.3390/vaccines10030467
31. Evans S, Klas A, Mikocka-Walus A, et al. "Poison" or "protection"? A mixed methods exploration of Australian parents' COVID-19 vaccination intentions. *J Psychosom Res* 2021;150:110626. doi:10.1016/j.jpsychores.2021.110626