

**Original article**

**DOI: 10.5633/amm.2024.0202**

**Longitudinal evaluation of the health-related quality of life in COVID-19 patients:  
a comparison between pre-infection and three months post-discharge**

Radomir Damjanović<sup>1</sup>, Iva Cvetković<sup>2</sup>, Maja Cvetanović<sup>2</sup>, Vuk Milošević<sup>1,3</sup>,

Aleksandar Stojanov<sup>1,3</sup>, Slobodan Vojinović<sup>1,3</sup>, Miodrag Vrbić<sup>2,3</sup>

<sup>1</sup>Clinic for Neurology, University Clinical Center of Niš, Niš, Serbia

<sup>2</sup>Clinic for Infectious Diseases, University Clinical Center of Niš, Niš, Serbia

<sup>3</sup>Faculty of Medicine, University of Niš, Niš, Serbia

Contact: Radomir Damjanović

e-mail: rdamjanovic@gmail.com

Dr. Zoran Djindjic Blvd., 18000 Niš, Serbia

**Originalni rad**

**DOI: 10.5633/amm.2024.0202**

**Procena kvaliteta života povezana sa zdravljem kod pacijenata sa COVID-19:  
upoređivanje perioda pre infekcije i tri meseca nakon otpusta sa bolničkog lečenja**

Radomir Damjanović<sup>1</sup>, Iva Cvetković<sup>2</sup>, Maja Cvetanović<sup>2</sup>, Vuk Milošević<sup>1,3</sup>,

Aleksandar Stojanov<sup>1,3</sup>, Slobodan Vojinović<sup>1,3</sup>, Miodrag Vrbić<sup>2,3</sup>

<sup>1</sup>Klinika za neurologiju, Univerzitetski klinički centar Niš, Republika Srbija

<sup>2</sup>Klinika za infektivne bolesti, Univerzitetski klinički centar Niš, Republika Srbija

<sup>3</sup>Medicinski fakultet, Univerzitet u Nišu, Republika Srbija

Kontakt: Radomir Damjanović

e-mail: rdamjanovic@gmail.com

Bul. dr Zorana Đinđića, 48, 18000 Niš, Srbija

## Sažetak

**Uvod:** Infektivno oboljenje Covid-19, izazvano SARS-CoV-2 virusom, i dalje predstavlja značajnu globalnu pretnju po zdravlje ljudi. Pored nedvosmislenih efekata na fizičko, postoje dokazi o ozbiljnom uticaju ovog oboljenja i na mentalno zdravlje.

**Cilj:** Naše istraživanje je imalo za cilj da proceni promene u kvalitetu života kod pacijenata obolelih od COVID-19 pre hospitalizacije i tri meseca nakon otpusta.

**Metodologija:** Podaci su dobijeni od 70 ispitanika sa postavljenom dijagnozom COVID-19, hospitalizovanih na Klinici za infektivne bolesti UKC Niš. Oni su ispitani dva puta, nakon prijema na bolničko lečenje i tri meseca nakon otpusta. Kvalitet života je meren korišćenjem sledećih upitnika: 36-item Short Form Survey (SF-36), Pittsburgh Sleep Quality Index (PSQI), Patient Health Questionnaire-9 (PHQ9) i Generalized Anxiety Disorder 7-item (GAD-7). Statistička značajnost dve promenljive u istom uzorku u dva različita vremenska trenutka je upoređivana korišćenjem uparenog T-testa.

**Rezultati:** Postojala je statistički značajna razlika u PHQ9 skorovima pre i nakon hospitalizacije ( $t = 4.738$ ,  $p < 0.01$ ). Nije nađena značajna promena PSQI skora pre hospitalizacije i tri meseca nakon otpusta ( $t = -.622$ ,  $p = .536$ ). Naši rezultati su pokazali statistički značajan porast GAD7 i značajan pad skora komponente fizičkog funkcionisanja (PF) ( $t = 5.929$ ,  $p < 0.01$ ) kao i skora komponente ograničenja zbog fizičkih zdravstvenih problema (RP) ( $t = 4.385$ ,  $p < 0.01$ ). Nalazi ukazuju da nije bilo statistički značajne razlike između perioda pre infekcije i tri meseca nakon otpusta u komponentama SF-36 upitnika koje se odnose na ograničenja usled emocionalnih problema (RE) i na domen socijalnog funkcionisanja (SF).

**Zaključak:** Rezultati našeg istraživanja ukazuju da pacijenti oboleli od COVID-19 imaju fizičke i mentalne zdravstvene probleme koji mogu trajati i nakon oporavka od infekcije i negativno uticati na njihov kvalitet života.

**Ključne reči:** kvalitet života, COVID-19, SF-36, PSQI, PHQ9, GAD-7

AMM Paper Accepted

**Longitudinal evaluation of the health-related quality of life in COVID-19 patients:  
a comparison between pre-infection and three months post-discharge**

**Abstract**

**Introduction:** The infectious disease COVID-19, caused by the SARS-CoV-2 virus, continues to be a significant global public health emergency. Alongside the undeniable effects on physical health, there is evidence of a serious impact on mental health as well.

**Aim:** Our study aimed to assess changes in health-related quality of life in COVID-19 patients before hospitalization for SARS-CoV-2 infection and three months after discharge.

**Methods:** Data were collected from 70 participants hospitalized for COVID-19. Participants were examined twice, after admission to the hospital and three months after discharge. Quality of life was measured with the 36-item Short Form Survey (SF-36), Pittsburgh Sleep Quality Index (PSQI), Patient Health Questionnaire-9 (PHQ9), and Generalized Anxiety Disorder 7-item (GAD-7). The statistical significance of two variables in the same sample at two distinct points in time was compared using a paired-sample T-test.

**Results:** There was a significant difference in PHQ9 scores before and after hospitalization ( $t = 4.738, p < 0.01$ ). We found no significant change in PSQI score before hospitalization or three months after discharge ( $t = -.622, p = .536$ ). Our results showed a significant increase in GAD7 and a significant decrease in physical functioning (PF) ( $t = 5.929, p < 0.01$ ) and role limitations due to physical health problems (RP) ( $t = 4.385, p < 0.01$ ) scores. The findings indicate that there was no statistically significant difference between pre-infection and three months after discharge in the SF-36 questionnaire's role limitations due to emotional issues (RE) and social functioning (SF) components.

**Conclusion:** In summary, the results of our study suggest that COVID-19 patients have physical and mental health problems that may persist even after recovery and negatively affect their quality of life.

**Keywords:** health-related quality of life, COVID-19, SF-36, PSQI, PHQ9, GAD-7

## **Introduction**

The infectious disease COVID-19, caused by the SARS-CoV-2 virus, continues to represent a significant global public health danger. The majority of patients who were released passed through various phases of recovery, while consequences on long-term health are presently being evaluated (1). Alongside the clear impacts on physical health, there is evidence of a substantial influence on mental health as well (2). Besides the symptoms of illness themselves, experiences during hospitalization, physical and social isolation, and death of family members or other patients are factors that could worsen mental health status (3). Additionally, coronavirus can induce immunological reactions that have serious effects on an infected person's mental health and brain function (4). The specific process by which coronavirus affects the brain is unclear. However, it can enter the brain via the olfactory bulb, pass the blood-brain barrier, and interact with ACE2 receptors (5). Furthermore, physiologic abnormalities in the brain, like microglia stimulation and cytokine activation may be connected to mental disorders (6). Experienced stress and compensating mechanisms are two well-known and plainly observable elements that effect one's actions and symptom presentations (7). Previous investigation on COVID-19 infection suggested that coronavirus infection was associated to long-term mental issues and neuropsychiatric effects (8). COVID-19 patients, in particular, exhibited symptoms of psychological distress symptoms that can result in persistent dysfunction and a reduction in their quality of life; recent data also show

an increased chance of having post-traumatic stress disorder during hospitalization (9, 10).

Here we provide a research in which health-related quality of life parameters were investigated in COVID-19 patients before and three months post hospital release.

## **Aim**

Our study aimed to assess changes in health-related quality of life in COVID-19 patients before hospitalization for SARS-CoV-2 infection and three months after discharge.

## **Methods**

### **Research Methodology and Participant Profile**

This observational, longitudinal study was carried out in the Infectious Diseases Clinic, University Clinical Center of Nis, Serbia, from January 2022 to May 2022. Participants were examined twice, during admission to the hospital and three months after discharge. Inclusion criteria were SARS-CoV-2 infection and a minimum age of 18. A positive PCR test done at the time of admission to our hospital proved the patient's infection with SARS-CoV-2. Participants were removed if they had poor questionnaire completion, exhibited evidence of prior mental illness or dementia, had spent more than 12 hours in the Intensive Care Unit, or experienced a fatal outcome. The participants' ages varied from 23 to 79. Demographic data and prior illnesses were gathered from hospital records and self-reported upon admission. Written informed permission was gained by each subject. The study followed the principles specified in the Helsinki Declaration. The University Clinical Center of Nis' Ethics Committee accorded their consent to the study proposal.

## Instruments

Quality of life was examined with the 36-item Short Form Survey (SF-36), Pittsburgh Sleep Quality Index (PSQI), Patient Health Questionnaire-9 (PHQ9), and Generalized Anxiety Disorder 7-item (GAD-7). A widely adopted measure for assessing health-related quality of life is the SF-36. Eight scales are measured by it: Physical functioning (PF) measures the ability to carry out physical tasks, including walking, using stairs, and house maintenance. Role limitations due to physical health problems (RP) measure the influence of physical health issues on everyday obligations, such as employment or other regular activities. Bodily pain (BP) measures the degree and effect of pain on everyday activities. The questionnaire covers items for general health (GH), vitality (VT), role limitations due to emotional issues (RE), social functioning (SF), and mental health (MH). General health (GH) examines self-reported overall health status, including perceived changes in health throughout time. Vitality (VT) examines the levels of energy and fatigue. Role limitations due to emotional issues (RE) analyzes the impact of emotional health problems on everyday activities, including employment or other regular activities. Mental health (MH) assesses psychological discomfort and well-being, whereas social functioning (SF) measures the effect of physical or emotional challenges on relationships (2). Scores vary from 0 to 100; lower scores imply greater dysfunction, while higher scores represent reduced dysfunction. The SF-36 analyzes two distinct dimensions: a physical dimension and a mental dimension. The assessment of the physical and mental factors is influenced by each scale in a different way (3). Instead of using an SF-36 overall score, for this study, we used the scores separately for PF, RP, SF, and RE in statistical processing. The Pittsburgh Sleep Quality Index, or PSQI, is a questionnaire developed to quantify a person's sleep quality over one month. Nineteen separate items make up the test, which examines seven major aspects of sleep quality: subjective sleep quality, sleep latency, sleep length, habitual sleep efficiency, disturbance in



sleep, usage of medications for sleep, and disruption of daily routines. One overall score is derived by adding the outcomes of these seven component scores (4). The Patient Health Questionnaire-9, known as PHQ-9, is a self-administered depression screening instrument. The Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) provides nine diagnostic criteria for major depressive disorder, which are reflected in the PHQ-9. With a total score that can vary from 0 to 27, each item is rated on a scale of 0 to 3. Higher scores suggest greater depression severity (5). The Generalized Anxiety Disorder-7-item Scale is referred to as the GAD-7. The existence and severity of generalized anxiety disorder (GAD) symptoms are assessed using this seven-item screening questionnaire (6).

### **Statistical Analysis**

Demographic data were presented as mean  $\pm$  SD. Descriptive statistical analysis was applied for the demographic information, and a paired-sample T-test was applied to evaluate the statistical significance of two variables in the same sample at two different points in time. IBM SPSS Statistics 26 was used for data processing, and  $p < 0.05$  was regarded as significant.

### **Results**

Data were gathered from 70 individuals hospitalized for COVID-19. Most of them were male (60%) and had a secondary level of education (45.71%). The mean age of the subjects was  $52.03 \pm 15.76$  years, and the median period since the positive PCR test was 4 days. Specific demographic information is presented in Table 1. The PHQ9 mean score calculated for the duration of hospitalization was  $6.600 \pm 4.284$ , and 3 months later it was  $4.314 \pm 3.0$ . In contrast, the PSQI and GAD-7 mean scores were higher three months following discharge. In addition, the mean values of RE and SF were similarly greater,

although PF and RP were decreased three months after infection (Table 2). To compare the scores of these questionnaires, we performed a paired-sample test for statistical significance between the scores obtained one month before COVID-19 infection and three months after hospital discharge (Table 3). PHQ9 scores before and after hospitalization differed significantly, according to our statistical findings ( $t = 4.738, p < 0.01$ ). We found no significant change in PSQI score before hospitalization and three months after discharge ( $t = -.622, p = .536$ ). According to our findings, the GAD7 score significantly increased 3 months after hospital discharge ( $t = -3.978, p < 0.01$ ), and there was a significant decline in physical functioning (PF) ( $t = 5.929, p < 0.01$ ) and role limitations due to physical health problems (RP) ( $t = 4.385, p < 0.01$ ) scores from pre-COVID-19 infection to 3 months post-hospital discharge. The results demonstrate that there was no difference in statistical significance in the role limitations due to emotional problems (RE) and social functioning (SF) components of the SF-36 questionnaire between pre-infection and 3 months after discharge. Paired samples t-test indicated a RE mean difference of  $-.043$  ( $SD = .428$ ), with a t-value of  $-.840$  and a p-value of  $.404$  and a SF mean difference of  $-.053$  ( $SD = .231$ ), with a t-value of  $-1.903$  and a p-value of  $.051$ .

**Table 1.** Demographic data of participants

<b>Variables</b>	<b>Total N (%), N=70</b>
Age in years (mean+SD)	52.03±15.76
Gender	
male	42 (60)
female	28 (40)
Education level	
Primary school	6 (8.57)
Secondary school	32 (45.71)
Faculty	24 (34.29)

Master degree	7 (10)
Doctoral degree	1 (1.43)
Marital status	
Single	9 (12.86)
Married	50 (71.43)
Divorced	6 (8.57)
Widowed	5 (7.14)
Employment status	
Student	3 (4.29)
Employed	40 (57.14)
Unemployed	27 (38.57)
Chronic illness	
Yes	41 (58.57)
No	29 (41.43)

**Table 2.** Descriptive statistics

Paired Samples Statistics					
		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	PHQ9	6.6000	70	4.28479	.51213
	PHQ9_3	4.3143	70	2.99537	.35802
Pair 2	PSQI	4.6000	70	3.16869	.37873
	PSQI_3	4.8000	70	2.59095	.30968
Pair 3	GAD7	5.3000	70	2.80966	.33582
	GAD7_3	7.1571	70	3.60610	.43101
Pair 4	PF	.8343	70	.15430	.01844
	PF_3	.6686	70	.28363	.03390
Pair 5	RP	.6500	70	.32804	.03921
	RP_3	.4464	70	.32385	.03871
Pair 6	RE	.7190	70	.31929	.03816
	RE_3	.7620	70	.28454	.03401
Pair 7	SF	.6632	70	.24923	.02979

SF\_3 .7157 70 .18945 .02264

**Table 3.** Paired Samples Correlations

		Paired Samples Test							
		Paired Differences					t	df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	PHQ9 - PHQ9_3	2.28571	4.03658	.48246	1.32323	3.24820	4.738	69	<b>.000</b>
Pair 2	PSQI - PSQI_3	-.20000	2.68976	.32149	-.84135	.44135	-.622	69	.536
Pair 3	GAD7 - GAD7_3	-1.8571	3.90572	.46682	-2.7884	-.92586	-3.978	69	<b>.000</b>
Pair 4	PF - PF_3	.16571	.23383	.02795	.10996	.22147	5.929	69	<b>.000</b>
Pair 5	RP - RP_3	.20357	.38845	.04643	.11095	.29619	4.385	69	<b>.000</b>
Pair 6	RE - RE_3	-.04300	.42820	.05118	-.14510	.05910	-.840	69	.404
Pair 7	SF - SF_3	-.05250	.23087	.02759	-.10755	.00255	-1.903	69	<b>.051</b>

p<0.01

## Discussion

As we are in the fourth year since the whole world was confronted with COVID-19, one of the most crucial problems facing global health systems today is to identify and comprehend the unavoidable effects of the SARS-CoV-2 viral infection.

Based on one of the first cross-sectional studies of individuals recovering from COVID-19 in Wuhan, SARS-CoV-2 virus infection is typically associated with several psychiatric disorders, including depression, anxiety, and even suicide. According to this study, 10% of COVID-19 survivors report anxiety or depression, and 29.5% of survivors experience sleep disturbances one month and six months after discharge, respectively (16). Our study's findings suggest that after hospital discharge, the features and standard of sleep recovered from COVID-19 do not dramatically alter. It is essential to keep in mind that the study's sample size was quite small, which could have reduced its ability to identify significant changes. Only a few factors have been linked to improved sleep quality. Poor sleep quality has been connected to female sex, younger age, and persons with a history of mental issues (17). Lack of sleep severely influences cognitive function and quality of life (18, 19). In addition, sleep interruptions could increase depression and anxiety while reducing pain thresholds (20).

A recently published meta-analysis analyzed the cumulative prevalence of mental diseases among COVID-19 survivors (21). Despite significant variance across the 27 studies, it was clear that mental and psychological health outcomes are essential for COVID-19 survivors. Prevalence rates for posttraumatic stress disorder (PTSD), anxiety, psychological distress, depression, and sleep disorders were as follows: 22% for anxiety, 36% for psychological distress, 21% for depression, and 35% for sleep disorders. The following risk variables have been connected to an enhanced risk of anxiety, depression, or other unfavorable mental health outcomes: indicators of inflammation, especially IL-6 (22), severity of illness, duration of symptoms, severity of illness at 6-month follow-up, and female

sex (23,24). Based on our results, there seems to be a statistically significant difference in the incidence of generalized anxiety disorder before and after COVID -19 infection, as shown by the GAD7 questionnaire. It is noteworthy to underline that the elevation in GAD7 scores identified in our study could be connected to a range of factors, such as the physical and emotional stress associated with hospitalization and treatment for COVID-19, as well as the social and economic disruption generated by the pandemic (25,26). One Chinese study applied the SF-36 questionnaire to investigate the health-related quality of life of COVID-19 patients (27). This showed a worse score at one-month follow-up, with psychological impairment described mainly in women, suggesting that the female gender may be a risk factor for psychological quality of life in these patients (27). In our study, the Physical Functioning (PF) component of the SF-36 was compared 3 months after discharge to the period before infection. The results showed a significant decrease in the ability to perform physical activities after the patients had recovered from COVID-19. These findings support other studies that have shown COVID-19 to have long-term health impacts, including reduced physical functioning (1). Our findings also indicate a significant decline in role limitations due to physical health problems (RP) - as a measure of the impact of physical health problems on daily activities among those who had COVID-19 and recovered. The reduction in physical role functioning may have several effects on the overall well-being of COVID-19 survivors. For instance, it would make it more difficult for them to carry out routine tasks like work or household, which would increase stress and reduce life pleasure (27). Another Chinese study that evaluated health-related quality of life using the SF-36 found that COVID-19 hospitalization had a detrimental impact on patients' quality of life for up to three months following release (12). In this research, poor quality of life appeared to be related to age, gender, and physical symptoms after discharge, suggesting that older adults,

particularly women and those with recurrent physical symptoms, are most at risk for poor quality of life.

### **Conclusion**

In summary, the results of our study demonstrate that COVID-19 patients have physical and mental health issues that may persist even after recovery and negatively affect their quality of life. The fact that our study was done at one site and the sample size was quite limited should be emphasized, as this may hamper the generalizability of our findings. However, our study underlines the importance of continuing surveillance and care for COVID-19 survivors to make sure they receive the necessary treatment to address any long-term health concerns. The long-term health implications of COVID-19 may be better understood in the future through research utilizing broader samples and more diverse populations.

## References

1. Huang C, Huang L, Wang Y et al. 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *Lancet*. 2021;397(10270):220–32. doi:10.1016/S0140-6736(20)32656-8
2. Li D., Liao X., Ma Z. *et al.* Clinical status of patients 1 year after hospital discharge following recovery from COVID-19: a prospective cohort study. *Ann. Intensive Care* 2022; **12**; 64. <https://doi.org/10.1186/s13613-022-01034-4>
3. Liu D, Baumeister RF, Veilleux JC, et al. Risk factors associated with mental illness in hospital discharged patients infected with COVID-19 in Wuhan, China. *Psychiatry Res*. 2020;292:113297. doi:10.1016/j.psychres.2020.113297
4. Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry*. 2020;7(6):547-560. doi:10.1016/S2215-0366(20)30168-1.
5. Gupta PK, Singh S, Mahour P, et al. Mental health outcome in hospitalized COVID-19 patients: An observational analysis from North Indian tertiary care hospital. *Clin Epidemiol Glob Health*. 2023;19:101209. doi:10.1016/j.cegh.2022.101209
6. Szcześniak D, Gładka A, Misiak B, Cyran A, Rymaszewska J. The SARS-CoV-2 and mental health: From biological mechanisms to social consequences. *Prog*



7. Kar N, Kar B, Kar S. Stress and coping during COVID-19 pandemic: Result of an online survey. *Psychiatry Res.* 2021;295:113598. doi:10.1016/j.psychres.2020.113598
8. Lam MH, Wing YK, Yu MW, et al. Mental morbidities and chronic fatigue in severe acute respiratory syndrome survivors: long-term follow-up. *Arch Intern Med.* 2009;169(22):2142-2147. doi:10.1001/archinternmed.2009.384
9. Tarsitani L, Vassalini P, Koukopoulos A, et al. Post-traumatic Stress Disorder Among COVID-19 Survivors at 3-Month Follow-up After Hospital Discharge. *J Gen Intern Med.* 2021;36(6):1702-1707. doi:10.1007/s11606-021-06731-7
10. Rogers JP, Chesney E, Oliver D, et al. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry.* 2020;7(7):611-627. doi:10.1016/S2215-0366(20)30203-0
11. McHorney CA, Ware JE Jr, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care.* 1993;31(3):247-263. doi:10.1097/00005650-199303000-00006
12. Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE Open Med.* 2016;4;4:2050312116671725. doi:10.1177/2050312116671725
13. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989 May;28(2):193-213. doi:10.1016/0165-1781(89)90047-4
14. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606-613. doi:10.1046/j.1525-1497.2001.016009606.x
15. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: The GAD-7. *Arch Intern Med.* 2006 May 22;166(10): 1092-7. doi:10.1001/archinte.166.10.1092.

16. Wu C, Hu X, Song J, et al. Mental health status and related influencing factors of COVID-19 survivors in Wuhan, China. *Clin Transl Med.* 2020;10(2):e52. doi:10.1002/ctm2.52
17. Mazza MG, De Lorenzo R, Conte C, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behav Immun.* 2020;89:594-600. doi:10.1016/j.bbi.2020.07.037
18. Chandra SS, Loganathan K, Awuzie BO, Wang F. A Longitudinal Study Examining the Association between Cognitive Behavior and Rational Abilities and the Effect of Sleep Quality on Construction Laborers. *Sustainability.* 2023; 15(7):6257. <https://doi.org/10.3390/su15076257>
19. Şahin, H.; Yıldırım, A.; Hacıhasanoğlu Aşlar, R.; Çebi, K.; Güneş, D. The relationship between nutritional behaviours and sleep quality in individuals applying to primary healthcare organizations. *J. Turk. Sleep Med.* 2020, 7, 29–39.
20. Wei Y, Blanken TF, Van Someren EJW. Insomnia really hurts: effect of a bad night's sleep on pain increases with insomnia severity. *Front Psychiatry.* 2019;9:377. doi:10.3389/fpsy.2018.00377
21. Khraisat B, Toubasi A, AlZoubi L, Al-Sayegh T, Mansour A. Meta-analysis of prevalence: the psychological sequelae among COVID-19 survivors. *Int J Psychiatry Clin Pract.* (2021) 28:1–10. doi: 10.1080/13651501.2021.1993924
22. Kappelmann N, Dantzer R, Khandaker GM. Interleukin-6 as potential mediator of long-term neuropsychiatric symptoms of COVID-19. *Psychoneuroendocrinology.* 2021;131:105295. doi:10.1016/j.psyneuen.2021.105295
23. Putri C, Arisa J, Hananto JE, Hariyanto TI, Kurniawan A. Psychiatric sequelae in COVID-19 survivors: A narrative review. *World J Psychiatry.* 2021;11(10):821-829. doi:10.5498/wjp.v11.i10.821
24. Schou TM, Joca S, Wegener G, Bay-Richter C. Psychiatric and neuropsychiatric sequelae of COVID-19 – A systematic review. *Brain Behav Immun.* (2021) 97:328–48. doi: 10.1016/j.bbi.2021.07.018

25. Xiong J, Lipsitz O, Nasri F, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord.* 2020;277:55-64. doi:10.1016/j.jad.2020.08.001
26. Wang C, Pan R, Wan X, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun.* 2020;87:40-48. doi:10.1016/j.bbi.2020.04.028
27. Chen KY, Li T, Gong FH, Zhang JS, Li XK. Predictors of Health-Related Quality of Life and Influencing Factors for COVID-19 Patients, a Follow-Up at One Month. *Front Psychiatry.* 2020;11:668. Published 2020 Jul 8. doi:10.3389/fpsy.2020.00668

AMM Paper Accepted