

## INTERSTITIAL PREGNANCY AFTER IPSILATERAL SALPINGECTOMY - CASE REPORT

*Žaklina Savić Mitić<sup>1</sup>, Mirko Trailović<sup>1</sup>, Dragiša Mitić<sup>2</sup>*

Health Centre Zaječar, Department of Gynecology and Obstetrics, Zaječar, Serbia<sup>1</sup>  
Public Health Institution "Timok", Zaječar, Serbia<sup>2</sup>

Contact: Žaklina Savić Mitić  
Požarevačka 63/7, 19000 Zaječar, Serbia  
E-mail: zaklina1009@gmail.com

Interstitial pregnancy is ectopic pregnancy which occurs within the intramural (interstitial) part of the uterine tube. The frequency of its occurrence is 3-4% of all ectopic pregnancies. It is also significant because of a long asymptomatic period and common ending with the rupture of the uterine horn as a serious complication. Mortality after this type of ectopic pregnancy is about 2,5%. This paper presents a case of a 29-year-old patient hospitalized for suspected ectopic interstitial pregnancy in the left uterine horn. Her anamnesis showed the left tube pregnancy ended with salpingectomy after the rupture of the uterine tube. The patient was examined both clinically and by ultrasound, her  $\beta$ -HCG level was being monitored, and after an adequate preoperative preparation, she underwent a surgery. Surgical intervention consisted of the incision of pregnancy-altered uterine horn, curettage and evacuation of ovulatory tissue, which saved the uterine horn. Timely diagnosis of interstitial pregnancy can prevent its possible serious complications. *Acta Medica Medianae 2015;54(2):52-55.*

**Key words:** *interstitial pregnancy, ectopic pregnancy, salpingectomy*