

## EXTRAMAMMARY PAGET'S DISEASE OF THE INGUINUM: A CASE REPORT

Vesna Karanikolić<sup>1,4</sup>, Aleksandar Karanikolić<sup>2,4</sup>, Dejan Petrović<sup>3,4</sup>, Danijela Popović<sup>1</sup>,  
Maša Golubović<sup>1</sup>, Miodrag Djordjević<sup>2</sup>

<sup>1</sup>Clinic of dermatology, Clinical center Niš, Serbia

<sup>2</sup>Surgical clinic, Clinical center Niš, Serbia

<sup>3</sup>Institute for cardiovascular disease Niška Banja, Serbia

<sup>4</sup>University of Niš, Faculty of Medicine, Niš, Serbia

Contact: Vesna Karanikolić  
Vardarska 23, 18000 Niš, Serbia  
E-mail: pean@ptt.rs

Extramammary Paget's disease (EMPD) is a rare intra-epithelial malignancy that is occasionally associated with an invasive adenocarcinoma component as well as other secondary cancers. The diagnosis is confirmed by the presence of Paget's cells on histopathological examination of a tissue specimen. The standard treatment method of inguinal EMPD, as well as EMPD in other areas, is surgical resection. We present a patient with extra-mammary Paget's disease, who developed the erythematous eruption in the right inguinal region.

*Acta Medica Medianae 2018;57(2):31-33.*

**Key words:** Extramammary Paget's disease, inguinum, diagnosis, treatment